

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317335876>

# Comprehensive compilation of randomized controlled trials (RCTs) involving psychodynamic treatments and interventions

Research · June 2017

CITATIONS

0

READS

1,664

1 author:



[Peter Lillengren](#)

Ersta Sköndal Bräcke University College, Stockholm, Sweden

21 PUBLICATIONS 100 CITATIONS

[SEE PROFILE](#)

# Comprehensive compilation of randomized controlled trials (RCTs) involving psychodynamic treatments and interventions

Peter Lilliengren, PhD  
Ersta Sköndal Bräcke University College  
Stockholm, Sweden  
[peter.lilliengren@esh.se](mailto:peter.lilliengren@esh.se)

Last updated: 2018-12-27

#	Reference (primary and secondary articles)	Patient sample	PDT-model	Comparison group(s)
1.	<b>Abbass, A., Sheldon, A., Gyra, J., &amp; Kalpin, A.</b> (2008). Intensive short-term dynamic psychotherapy for DSM-IV personality disorders: a randomized controlled trial. <i>J Nerv Ment Dis</i> , 196, 211-216. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18340256">http://www.ncbi.nlm.nih.gov/pubmed/18340256</a>	<b>Personality disorders</b> (mixed)	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Davanloo, 1990)	<b>Minimal contact</b>
2.	<b>Ahola, P., Joensuu, M., Knekt, P., Lindfors, O., Saarinen, P., Tolmunen, T., ... Lehtonen, J.</b> (2016). Effects of Scheduled Waiting for Psychotherapy in Patients With Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 0(0), 1. <a href="http://doi.org/10.1097/NMD.0000000000000616">http://doi.org/10.1097/NMD.0000000000000616</a>	<b>Depression</b>	<b>Long-Term Psychodynamic Psychotherapy</b> (At least 12 months, biweekly sessions, no reference to manual)	<b>Waitlist</b> (6 months)
3.	<b>Ajlilchi, B., Nejati, V., Town, J. M., Wilson, R., &amp; Abbass, A.</b> (2016). Effects of Intensive Short-Term Dynamic Psychotherapy on depressive symptoms and executive functioning in Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 204(7), 500–5. <a href="http://doi.org/10.1097/NMD.0000000000000518">http://doi.org/10.1097/NMD.0000000000000518</a>	<b>Depression</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Davanloo, 1990)	<b>Waitlist</b>
4.	<b>Alstrom, J. E., Nordlund, C. L., Persson, G., Harding, M., &amp; Ljungqvist, C.</b> (1984). Effects of four treatment methods on social phobic patients not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 97–110. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6148843">http://www.ncbi.nlm.nih.gov/pubmed/6148843</a>	<b>Social anxiety disorder (SAD)</b>	<b>Supportive dynamic psychotherapy</b> (Dewald, 1964)	<b>Prolonged exposure</b> (in vivo w. therapist)/ + <b>relaxation therapy</b>
5.	<b>Alstrom, J. E., Nordlund, G., Persson, G., Harding, M., &amp; Lundquist, C.</b> (1984). Effects of four treatment methods on agoraphobic women not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 1–17. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6147066">http://www.ncbi.nlm.nih.gov/pubmed/6147066</a>	<b>Agoraphobia</b> (in women not suitable for insight-oriented psychotherapy)	<b>Supportive dynamic psychotherapy</b> (Dewald, 1964)	<b>Prolonged exposure</b> (in vivo w. therapist)/ + <b>relaxation therapy</b>
6.	<b>Amianto, F., Ferrero, A., Pierò, A., Cairo, E., Rocca, G., Simonelli, B., ... Fassino, S.</b> (2011). Supervised team management, with or without structured psychotherapy, in heavy users of a mental health service with borderline personality disorder: a two-year follow-up preliminary randomized study. <i>BMC psychiatry</i> , 11(1), 181. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260092&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260092&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Borderline personality disorder</b>	<b>Sequential Brief Adlerian Psychodynamic Psychotherapy</b> (SB-APP; Fassino, 2008; 2009)	<b>Supervised team management</b>

7.	<p><b>Andersson, G., Paxling, B., Roch-Norlund, P., Ostman, G., Norgren, A., Almlöv, J., ... Silverberg, F. (2012).</b> Internet-Based Psychodynamic versus Cognitive Behavioral Guided Self-Help for Generalized Anxiety Disorder: A Randomized Controlled Trial. <i>Psychotherapy and psychosomatics</i>, 81(6), 344–355.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/22964540">http://www.ncbi.nlm.nih.gov/pubmed/22964540</a></p>	<b>Generalized Anxiety Disorder (GAD)</b>	<b>IPDT (SUBGAP; Silverberg, 2005)</b>	<b>ICBT (Paxling, 2011)</b>
8.	<p><b>Arnevik, E., Wilberg, T., Urnes, O., Johansen, M., Monsen, J. T., &amp; Karterud, S. (2009).</b> Psychotherapy for personality disorders: short-term day hospital psychotherapy versus outpatient individual therapy - a randomized controlled study. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 24(2), 71–8.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/19097870">http://www.ncbi.nlm.nih.gov/pubmed/19097870</a></p> <p>Arnevik, E., Wilberg, T., Urnes, Ø., Johansen, M., Monsen, J. T., &amp; Karterud, S. (2010). Psychotherapy for Personality Disorders: 18 Months' Follow-Up of the Ullevål Personality Project. <i>Journal of Personality Disorders</i>, 24(2), 188–203.  <a href="http://guilfordjournals.com/doi/pdf/10.1521/pedi.2010.24.2.188">http://guilfordjournals.com/doi/pdf/10.1521/pedi.2010.24.2.188</a></p>	<b>Personality disorders (mixed)</b>	<b>Short-Term Day Hospital Psychotherapy (DHP; no specific reference)</b>	<b>Outpatient Individual Psychotherapy</b>
9.	<p><b>Bachar, E., Latzer, Y., Kreitler, S., &amp; Berry, E. M. (1999).</b> Empirical comparison of two psychological therapies. Self psychology and cognitive orientation in the treatment of anorexia and bulimia. <i>J Psychother Pract Res</i>, 8, 115-128.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/10079459">http://www.ncbi.nlm.nih.gov/pubmed/10079459</a></p>	<b>Bulimia</b>	<b>Self Psychological Treatment (based on Kohut, 1984)</b>	<b>Cognitive Orientation Treatment (Kreitler &amp; Kreitler, 1990)</b>
10.	<p><b>Baldoni, F., Baldaro, B., &amp; Trombini, G. (1995).</b> Psychotherapeutic perspectives in urethral syndrome. <i>Stress Medicine</i>, 11(July 1994), 79–84.  <a href="http://onlinelibrary.wiley.com/doi/10.1002/smi.2460110115/abstract">http://onlinelibrary.wiley.com/doi/10.1002/smi.2460110115/abstract</a></p>	<b>Urethral syndrome</b>	<b>STDP (Malan, 1979; Davanloo, 1980)</b>	<b>TAU</b>
11.	<p><b>Balottin, U., Ferri, M., Racca, M., Rossi, M., Rossi, G., Beghi, E., ... Termine, C. (2014).</b> Psychotherapy versus usual care in pediatric migraine and tension-type headache: a single-blind controlled pilot study. <i>Italian Journal of Pediatrics</i>, 40(1), 6.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/24444140">http://www.ncbi.nlm.nih.gov/pubmed/24444140</a></p>	<b>Idiopathic headache (children/adolescents 6-18 yrs)</b>	<b>STPP (Aliprandi et al, 1990)</b>	<b>TAU</b>
12.	<p><b>Barber, J. P., Barrett, M. S., Gallop, R., Rynn, M. a, &amp; Rickels, K. (2011).</b> Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. <i>The Journal of clinical psychiatry</i>, 1–8.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/22152401">http://www.ncbi.nlm.nih.gov/pubmed/22152401</a></p> <p>Zilcha-Mano, S., Dinger, U., McCarthy, K. S., Barrett, M. S., &amp; Barber, J. P. (2013). Changes in well-being and quality of life in a randomized trial comparing dynamic psychotherapy and pharmacotherapy for</p>	<b>Depression</b>	<b>Supportive Expressive Therapy (Luborsky, 1984)</b>	<b>1. Placebo</b> <b>2. SSRI/SNRI</b>

	major depressive disorder. <i>Journal of affective disorders</i> , 1–5. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24176534">http://www.ncbi.nlm.nih.gov/pubmed/24176534</a>			
13.	<b>Barkham, M., Rees, A., Shapiro, D. A., Stiles, W. B., Agnew, R. M., &amp; Halstead, J., et al. (1996).</b> Outcomes of time-limited psychotherapy in applied settings: replicating the Second Sheffield Psychotherapy Project. <i>J Consult Clin Psychol</i> , 64, 1079-1085. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8916639">http://www.ncbi.nlm.nih.gov/pubmed/8916639</a>	<b>Depression</b>	<b>Psychodynamic-interpersonal Therapy</b> (Hobson, 1985)	<b>CBT</b> (no reference)
14.	<b>Barkham, M., Shapiro, D. A., Hardy, G. E., &amp; Rees, A. (1999).</b> Psychotherapy in two-plus-one sessions: outcomes of a randomized controlled trial of cognitive-behavioral and psychodynamic-interpersonal therapy for subsyndromal depression. <i>J Consult Clin Psychol</i> , 67, 201-211. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10224730">http://www.ncbi.nlm.nih.gov/pubmed/10224730</a>	<b>Depression</b> (subsyndromal)	<b>Psychodynamic-interpersonal Therapy</b> (3 sessions; Barkham & Shapiro, based on Hobson, 1985)	<b>CBT</b> (Beck et al 1979; Beck, 1995)
15.	<b>Bassett, D. L., &amp; Pilowsky, I. (1985).</b> A study of brief psychotherapy for chronic pain. <i>J Psychosom Res</i> , 29, 259-264. <a href="http://www.sciencedirect.com/science/article/pii/0022399885900522">http://www.sciencedirect.com/science/article/pii/0022399885900522</a>	<b>Chronic pain</b>	<b>12 sessions PDT</b> (no reference)	<b>Supportive psychotherapy</b> (no reference)
16.	<b>Bastos, A. G., Guimaraes, L. S. P., &amp; Trentini, C. M. (2014).</b> The efficacy of long-term psychodynamic psychotherapy, fluoxetine and their combination in the outpatient treatment of depression. <i>Psychotherapy Research : Journal of the Society for Psychotherapy Research</i> , (July 2014), 1–13. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25041333">http://www.ncbi.nlm.nih.gov/pubmed/25041333</a>  <b>Bastos, A. G., Pinto Guimarães, L. S., &amp; Trentini, C. M. (2013).</b> Neurocognitive changes in depressed patients in psychodynamic psychotherapy, therapy with fluoxetine and combination therapy. <i>Journal of Affective Disorders</i> , 1–10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24103853">http://www.ncbi.nlm.nih.gov/pubmed/24103853</a>	<b>Depression</b>	<b>Long-term psychodynamic psychotherapy</b> (LTPP; Gabbard, 2004; 2010)	<b>1. Fluoxetine alone</b> <b>2. LTPP + fluoxetine</b>
17.	<b>Bateman, A., &amp; Fonagy, P. (1999).</b> Effectiveness of partial hospitalization in the treatment of borderline personality disorder: a randomized controlled trial. <i>The American Journal of Psychiatry</i> , 156(10), 1563–9. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10518167">http://www.ncbi.nlm.nih.gov/pubmed/10518167</a>  <b>Bateman, A., &amp; Fonagy, P. (2008).</b> 8-Year Follow-Up of Patients Treated for Borderline Personality Disorder: Mentalization-Based Treatment Versus Treatment As Usual. <i>The American Journal of Psychiatry</i> , 165(5), 631–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18347003">http://www.ncbi.nlm.nih.gov/pubmed/18347003</a>	<b>Borderline personality disorder</b>	<b>Mentalization based treatment</b> (MBT; Bateman & Fonagy, 2004)	<b>TAU</b>
18.	<b>Bateman, A., &amp; Fonagy, P. (2009).</b> Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1355-1364. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19833">http://www.ncbi.nlm.nih.gov/pubmed/19833</a>	<b>Borderline personality disorder</b>  <b>Borderline and co-morbid Antisocial personality disorder</b>	<b>Mentalization based treatment</b> (MBT; Bateman & Fonagy, 2004)	<b>Structured Clinical Management (SCM)</b>

	<a href="#">787</a> Bateman, A., O'Connell, J., Lorenzini, N., Gardner, T., & Fonagy, P. (2016). A randomised controlled trial of mentalization-based treatment versus structured clinical management for patients with comorbid borderline personality disorder and antisocial personality disorder. <i>BMC Psychiatry</i> , 16(1), 304. <a href="http://bmcp psychiatry.biomedcentral.com/articles/10.1186/s12888-016-1000-9">http://bmcp psychiatry.biomedcentral.com/articles/10.1186/s12888-016-1000-9</a>	(subsample in secondary study)		
19.	<b>Bateman, A., Fonagy, P., Bateman, A., &amp; Fonagy, P.</b> (2018). A Randomized Controlled Trial of a Mentalization-Based Intervention (MBT-FACTS) for Families of People With Borderline Personality Disorder. <i>Personality Disorders: Theory, Research, and Treatment</i> .	<b>Families of people with Borderline Personality Disorder</b>	<b>MBT-FACTS</b> (psychoeducation and skills-training based on MBT; Bateman & Fonagy, 2016)	<b>Waitlist</b>
20.	<b>Ben-Itzhak, S., Bluvstein, I., Schreiber, S., Aharonov-Zaig, I., Maor, M., Lipnik, R., &amp; Bloch, M.</b> (2012). The Effectiveness of Brief Versus Intermediate Duration Psychodynamic Psychotherapy in the Treatment of Adjustment Disorder. <i>Journal of Contemporary Psychotherapy</i> , 42(4), 249–256. <a href="http://www.springerlink.com/index/10.1007/s10879-012-9208-6">http://www.springerlink.com/index/10.1007/s10879-012-9208-6</a>	<b>Adjustment disorder</b>	<b>Brief Psychodynamic Psychotherapy</b> (12 sessions; Malan, 1979)	<b>Brief Psychodynamic Psychotherapy</b> (1 yrs; Malan, 1979)
21.	<b>Beresnevaite, M.</b> (2000) Exploring the benefits of group psychotherapy in reducing alexithymia in coronary heart disease patients: a preliminary study. <i>Psychotherapy and Psychosomatics</i> , 69, 117–122. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10773774?dopt=Abstract">http://www.ncbi.nlm.nih.gov/pubmed/10773774?dopt=Abstract</a>	<b>Alexithymia</b> (in patients with coronary heart disease)	<b>Group psychotherapy</b> (no reference)	<b>Psychoeducation</b>
22.	<b>Bernik, M., Corregiari, F., Savoia, M. G., Barros Neto, T. P. de, Pinheiro, C., &amp; Neto, F. L.</b> (2018). Concomitant treatment with sertraline and social skills training improves social skills acquisition in social anxiety disorder: A double-blind, randomized controlled trial. <i>PLOS ONE</i> , 13(10), 2–18. <a href="http://doi.org/10.1371/journal.pone.0205809">http://doi.org/10.1371/journal.pone.0205809</a>	<b>Social anxiety disorder (SAD)</b>	<b>Group psychodynamic therapy</b> (GPT; no manual)  1. Sertraline + GPT  2. Placebo + GPT	<b>Group CBT</b> (manualized, but no reference)  1. Sertraline + GCBT  2. Placebo + GCBT
23.	<b>Beutel, M. E., Dippel, A., Szczepanski, M., Thiede, R., &amp; Wiltink, J.</b> (2006). Mid-Term Effectiveness of Behavioral and Psychodynamic Inpatient Treatments of Severe Obesity Based on a Randomized Study. <i>Psychotherapy and Psychosomatics</i> , 75(6), 337–345. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17053334?dopt=Abstract">http://www.ncbi.nlm.nih.gov/pubmed/17053334?dopt=Abstract</a>	<b>Severe obesity</b>	<b>Individual in-patient PDT</b> (Local program, no reference)	<b>In-patient CBT</b> (Local program, no reference)
24.	<b>Beutel, M. E., Scheurich, V., Knebel, A., Michal, M., Wiltink, J., Graf-Morgenstern, M., ... Subic-Wrana, C.</b> (2013). Implementing panic-focused psychodynamic psychotherapy into clinical practice. <i>Canadian journal of psychiatry. Revue canadienne de psychiatrie</i> , 58(6), 326–34. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23768260">http://www.ncbi.nlm.nih.gov/pubmed/23768260</a>	<b>Panic disorder</b>	<b>Panic Focused Psychodynamic Psychotherapy</b> (PFPP; Milrod)	<b>CBT</b>

25.	<p>Beutel, M. E., Weissflog, G., Leuteritz, K., Wiltink, J., Haselbacher, a., Ruckes, C., ... Brahler, E. (2013). Efficacy of short-term psychodynamic psychotherapy (STPP) with depressed breast cancer patients: results of a randomized controlled multicenter trial. <i>Annals of Oncology</i>, 1–7. <a href="http://annonc.oxfordjournals.org/cgi/doi/10.1093/annonc/mdt526">http://annonc.oxfordjournals.org/cgi/doi/10.1093/annonc/mdt526</a></p> <p>Weißflog, G., Brähler, E., Leuteritz, K., Barthel, Y., Kuhnt, S., Wiltink, J., ... Beutel, M. E. (2015). Does psychodynamic short-term psychotherapy for depressed breast cancer patients also improve fatigue? Results from a randomized controlled trial. <i>Breast Cancer Research and Treatment</i>. <a href="http://link.springer.com/10.1007/s10549-015-3494-0">http://link.springer.com/10.1007/s10549-015-3494-0</a></p>	Depression (in breast cancer patients)	Supportive-expressive therapy (Haselbacher, Barthel, Brähler et al., 2010; based on Luborsky, 1984)	TAU
26.	<p>Bloch, M., Meiboom, H., Lorberblatt, M., Bluvstein, I., Aharonov, I., &amp; Schreiber, S. (2012). The effect of sertraline add-on to brief dynamic psychotherapy for the treatment of postpartum depression: a randomized, double-blind, placebo-controlled study. <i>The Journal of clinical psychiatry</i>, 73(2), 235–41. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22401479">http://www.ncbi.nlm.nih.gov/pubmed/22401479</a></p>	Depression (post-partum)	Brief Dynamic Psychotherapy (BDT; Malan, 1979) + placebo meds	Brief Dynamic Psychotherapy (BDT; Malan, 1979) + Sertraline
27.	<p>Bögels, S., Wijts, P., Oort, F., &amp; Sallaerts, S. J. (2014). Psychodynamic psychotherapy versus cognitive-behavior therapy for social anxiety disorder: An efficacy and partial effectiveness trial. <i>Depression and Anxiety</i>, 31(5), 363–373. <a href="http://doi.wiley.com/10.1002/da.22246">http://doi.wiley.com/10.1002/da.22246</a></p>	Social anxiety disorder (SAD)	STDP (Malan, 1979; also Luborsky 1984)	CBT (Clark & Wells, 1995)
28.	<p>Bressi, C., Porcellana, M., Marinaccio, P. M., Nocito, E. P., &amp; Magri, L. (2010). Short-term psychodynamic psychotherapy versus treatment as usual for depressive and anxiety disorders: a randomized clinical trial of efficacy. <i>The Journal of Nervous and Mental Disease</i>, 198(9), 647–52. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20823726">http://www.ncbi.nlm.nih.gov/pubmed/20823726</a></p>	Mixed sample (depression and anxiety)	Brief Dynamic Psychotherapy (BDT; Malan, 1976; Osimo & Malan, 1992)	TAU
29.	<p>Brockman, B., Poynton, A., Ryle, A., &amp; Watson, J. P. (1987). Effectiveness of time-limited therapy carried out by trainees. Comparison of two methods. <i>Br J Psychiatry</i>, 151, 602-610. <a href="http://bjp.rcpsych.org/content/151/5/602">http://bjp.rcpsych.org/content/151/5/602</a></p>	Mixed sample	Interpretive therapy (Mann & Goldman, 1982)	Cognitive-analytic therapy (CAT; Ryle, 1979; 1982)
30.	<p>Brodsky, H., &amp; Andrews, G. (1983). Brief psychotherapy in family practice. A controlled prospective intervention trial. <i>Br J Psychiatry</i>, 143, 11-19. <a href="http://bjp.rcpsych.org/content/143/1/11.short">http://bjp.rcpsych.org/content/143/1/11.short</a></p>	Mixed sample	STDP (Balant, Malan & Sifneos)	1. Family practitioner 2. No treatment
31.	<p>Brom, D., Kleber, R. J., &amp; Defares, P. B. (1989). Brief psychotherapy for posttraumatic stress disorders. <i>J Consult Clin Psychol</i>, 57, 607-612. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2571625">http://www.ncbi.nlm.nih.gov/pubmed/2571625</a></p>	PTSD	Brief Psychodynamic Therapy (Horowitz, 1976)	1. Trauma Desensitization (no specific reference) 2. Hypnotherapy (no specific reference)
32.	<p>Budman, S. H., Demby, A., Redondo, J. P., Hannan, M., Feldstein, M., &amp; Ring, J., et al. (1988). Comparative outcome in time-limited individual and group psychotherapy. <i>Int J Group Psychother</i>, 38, 63-86.</p>	Mixed sample	Time-limited individual psychotherapy (Mann, 1973)	Time-limited group psychotherapy (Budman & Bennett, 1983)

	<a href="http://www.tandfonline.com/doi/abs/10.1080/00207284.1988.11491085">http://www.tandfonline.com/doi/abs/10.1080/00207284.1988.11491085</a>			
33.	<b>Burnand, Y., Andreoli, A., Kolatte, E., Venturini, A., &amp; Rosset, N. (2002).</b> Psychodynamic psychotherapy and clomipramine in the treatment of major depression. <i>Psychiatr Serv</i> , 53, 585-590. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11986508">http://www.ncbi.nlm.nih.gov/pubmed/11986508</a>	<b>Depression</b>	<b>PDT + clomipramine,</b> (unclear which model, Safran & Muran is mentioned among references)	<b>Supportive care + clomipramine</b>
34.	<b>Burns, A., Guthrie, E., Marino-Francis, F., Busby, C., Morris, J., Russell, E., ... Byrne, J. (2005).</b> Brief psychotherapy in Alzheimer's disease. <i>British Journal Of Psychiatry</i> , 187(2), 143-147. <a href="http://bjp.rcpsych.org/content/187/2/143.short">http://bjp.rcpsych.org/content/187/2/143.short</a>	<b>Alzheimers disease</b>	<b>Psychodynamic-Interpersonal therapy</b> (Hobson, 1985)	<b>TAU</b>
35.	<b>Chavooshi, B., Mohammadkhani, P., &amp; Dolatshahee, B. (2016).</b> A Randomized Double-Blind Controlled Trial Comparing Davanloo's Intensive Short-Term Dynamic Psychotherapy as Internet-Delivered Versus Treatment as Usual for Medically Unexplained Pain: A Six-Month Pilot Study. <i>Psychosomatics</i> , 57(3), 292-300. <a href="http://linkinghub.elsevier.com/retrieve/pii/S033318216000025">http://linkinghub.elsevier.com/retrieve/pii/S033318216000025</a>	<b>Medically unexplained pain (MUP)</b>	<b>Internet-delivered Intensive Short-Term Dynamic Psychotherapy</b> (ID-ISTDP; Abbass, 2015)	<b>TAU</b>
36.	<b>Chavooshi, B., Mohammadkhani, P., &amp; Dolatshahee, B. (2016).</b> Efficacy of Intensive Short-Term Dynamic Psychotherapy for Medically Unexplained Pain: A Pilot Three-Armed Randomized Controlled Trial Comparison with Mindfulness-Based Stress Reduction. <i>Psychotherapy and Psychosomatics</i> , 123-125. <a href="http://www.karger.com/?doi=10.1159/000441698">http://www.karger.com/?doi=10.1159/000441698</a>	<b>Medically unexplained pain (MUP)</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Abbass, 2015)	<b>1. Mindfulness-based stress reduction (MBSR; no reference)</b> <b>2. TAU</b>
37.	<b>Chavooshi, B., Mohammadkhani, P., &amp; Dolatshahee, B. (2016).</b> Telemedicine vs. in-person delivery of intensive short-term dynamic psychotherapy for patients with medically unexplained pain: A 12-month randomized, controlled trial. <i>Journal of Telemedicine and Telecare</i> , 0(0), 1-9. <a href="http://jit.sagepub.com/lookup/doi/10.1177/1357633X15627382">http://jit.sagepub.com/lookup/doi/10.1177/1357633X15627382</a>	<b>Medically unexplained pain (MUP)</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Abbass, 2015)	<b>Internet-delivered ISTDP</b> (ID-ISTDP)
38.	<b>Chavooshi, B., Saberi, M., Tavallaie, S. A., &amp; Sahraei, H. (2017).</b> Psychotherapy for Medically Unexplained Pain: A Randomized Clinical Trial Comparing Intensive Short-Term Dynamic Psychotherapy and Cognitive Behavior Therapy. <i>Psychosomatics</i> , <a href="http://dx.doi.org/10.1016/j.psym.2017.01.003">http://dx.doi.org/10.1016/j.psym.2017.01.003</a>	<b>Medically unexplained pain (MUP)</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Abbass, 2015)	<b>CBT</b> (Turk et al., 1983)
39.	<b>Clarici, A., Pellizzoni, S., Guaschino, S., Alberico, S., Bembich, S., Giuliani, R., ... Panksepp, J. (2015).</b> Intranasal administration of oxytocin in postnatal depression: implications for psychodynamic psychotherapy from a randomized double-blind pilot study. <i>Frontiers in Psychology</i> , 06, 1-10. <a href="http://journal.frontiersin.org/article/10.3389/fpsyg.2015.00426/abstract">http://journal.frontiersin.org/article/10.3389/fpsyg.2015.00426/abstract</a>	<b>Depression (post-partum)</b>	<b>Psychodynamic approach</b> (non-manualized?) + <b>oxytocine</b>	<b>Psychodynamic approach</b> (non-manualized?) + <b>placebo</b>
40.	<b>Clarkin, J. F., Levy, K. N., Lenzenweger, M. F.,</b>	<b>Borderline personality</b>	<b>Transference-Focused</b>	<b>1. DBT</b> (Linehan, 1991)

	<p>&amp; Kernberg, O. F. (2007). Evaluating three treatments for borderline personality disorder: a multiwave study. <i>Am J Psychiatry</i>, 164, 922-928. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17541052">http://www.ncbi.nlm.nih.gov/pubmed/17541052</a></p> <p>Levy, K. N., Meehan, K. B., Kelly, K. M., Reynoso, J. S., Weber, M., Clarkin, J. F., &amp; Kernberg, O. F. (2006). Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. <i>Journal of Consulting and Clinical Psychology</i>, 74(6), 1027-40. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17154733">http://www.ncbi.nlm.nih.gov/pubmed/17154733</a></p>	disorder	Psychotherapy (TFP; Kernberg)	2. Supportive treatment (Rockland, 1992)
41.	<p><b>Connolly Gibbons, M. B., Thompson, S. M., Scott, K., Schauble, L. a, Mooney, T., Thompson, D., ... Crits-Christoph, P. (2012).</b> Supportive-expressive dynamic psychotherapy in the community mental health system: A pilot effectiveness trial for the treatment of depression. <i>Psychotherapy (Chicago, Ill.)</i>, 49(3), 303-16. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22962971">http://www.ncbi.nlm.nih.gov/pubmed/22962971</a></p>	Depression	Supportive-expressive therapy (manual adapted from Luborsky, 1984)	TAU
42.	<p><b>Connolly Gibbons, M. B., Gallop, R., Thompson, D., Luther, D., Crits-Christoph, K., Jacobs, J., ... Crits-Christoph, P. (2016).</b> Comparative effectiveness of Cognitive Therapy and Dynamic Psychotherapy for Major Depressive Disorder in a community mental health setting. <i>JAMA Psychiatry</i>, 19104, 1-8. <a href="http://doi.org/10.1001/jamapsychiatry.2016.1720">http://doi.org/10.1001/jamapsychiatry.2016.1720</a></p>	Depression	Supportive-expressive therapy (Book, 1998; Luborsky, 1984)	CBT (Beck, 1995)
43.	<p><b>Cooper, P. J., Murray, L., Wilson, A., &amp; Romaniuk, H. (2003).</b> Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. <i>Br J Psychiatry</i>, 182, 412-419. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12724244">http://www.ncbi.nlm.nih.gov/pubmed/12724244</a></p> <p>Murray, L., Cooper, P. J, Wilson, A., &amp; Romaniuk, H. (2003). Controlled trial of the short- and long-term effect of psychological treatment of post- partum depression: 2. Impact on the mother-child relationship and child outcome. <i>British Journal of Psychiatry</i>, 182, 420-7. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12724245">http://www.ncbi.nlm.nih.gov/pubmed/12724245</a></p>	Depression (post-partum)	Psychodynamic therapy (Cramer et al, 1990; Stern, 1995)	<p>1. TAU</p> <p>2. Non-directive counseling (Holden et al, 1989)</p> <p>3. Cognitive-Behavioural Therapy (Hawton et al, 1989)</p>
44.	<p><b>Covi, L., &amp; Lipman, R. S. (1987).</b> Cognitive behavioral group psychotherapy combined with imipramine in major depression. <i>Psychopharmacol Bull</i>, 23, 173-176. <a href="http://www.ncbi.nlm.nih.gov/pubmed/3602315">http://www.ncbi.nlm.nih.gov/pubmed/3602315</a></p>	Depression	Traditional Group Therapy (based on "psychodynamic-Interpersonal" theories but with no reference)	<p>1. CBT group therapy (Beck, 1979; Covi, 1982)</p> <p>2. CBT group therapy + Imipramine</p>
45.	<p><b>Covi, L., Lipman, R. S., Derogatis, L. R., Smith, J. E., &amp; Pattison, J. H. (1974).</b> Drugs and group psychotherapy in neurotic depression. <i>Am J Psychiatry</i>, 131, 191-198. <a href="http://ajp.psychiatryonline.org/doi/abs/10.11">http://ajp.psychiatryonline.org/doi/abs/10.11</a></p>	Depression (chronic)	Psychodynamically oriented group therapy (no reference) + medication or placebo	Brief supportive contact + medication or placebo



	<a href="http://ajp.131.2.191">76/ajp.131.2.191</a>			
46.	<p><b>Creed, F.,</b> Fernandes, L., Guthrie, E., Palmer, S., Ratcliffe, J., &amp; Read, N., et al. (2003). The cost-effectiveness of psychotherapy and paroxetine for severe irritable bowel syndrome. <i>Gastroenterology</i>, 124, 303-317. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12557136">http://www.ncbi.nlm.nih.gov/pubmed/12557136</a></p> <p>Hyphantis, T., Guthrie, E., Tomenson, B., &amp; Creed, F. (2009). Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome. <i>Pain</i>, 145(1-2), 196–203. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19643544">http://www.ncbi.nlm.nih.gov/pubmed/19643544</a></p>	<b>Irritable bowel syndrome</b>	<b>Psychodynamic-Interpersonal Therapy (PIT;</b> Hobson, 1985)	<b>TAU/SSRI</b>
47.	<p><b>Crits-Christoph, P.,</b> Gibbons, M. B. C., Narducci, J., Schamberger, M., &amp; Gallop, R. (2005). Interpersonal Problems and the Outcome of Interpersonally Oriented Psychodynamic Treatment of GAD. <i>Psychotherapy: Theory, Research, Practice, Training</i>, 42(2), 211–224. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.2.211">http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.2.211</a></p>	<b>Generalized anxiety disorder (GAD)</b>	<b>Supportive Expressive Therapy</b> (Luborsky (1984)	<b>Supportive therapy</b> (Borkovec & Matthews, 1988; Borkovec & Costello, 1993)
48.	<p><b>Crits-Christoph, P.,</b> Siqueland, L., Blaine, J., Frank, A., Luborsky, L., &amp; Onken, L. S., et al. (1999). Psychosocial treatments for cocaine dependence: National Institute on Drug Abuse Collaborative Cocaine Treatment Study. <i>Arch Gen Psychiatry</i>, 56, 493-502. <a href="http://archpsyc.jamanetwork.com/article.aspx?articleid=1673777">http://archpsyc.jamanetwork.com/article.aspx?articleid=1673777</a></p> <p>Crits-Christoph, P., Gibbons, M. B. C., Gallop, R., Ring-Kurtz, S., Barber, J. P., Worley, M., ... Hearon, B. (2008). Supportive-Expressive Psychodynamic Therapy for Cocaine Dependence: A Closer Look. <i>Psychoanalytic Psychology : The Official Journal of the Division of Psychoanalysis, American Psychological Association, Division 39</i>, 25(3), 483–498. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2786223&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2786223&amp;tool=pmcentrez&amp;rendertype=abstract</a></p>	<b>Cocaine dependence</b>	<b>Supportive Expressive Therapy</b> (Luborsky (1984) + <b>group drug counseling</b> (Mercer & Woody, 1992)	<p><b>1. Cognitive therapy</b> (Beck, 1993) + <b>group drug counseling</b> (Mercer &amp; Woody, 1992)</p> <p><b>2. Group drug counseling alone</b></p>
49.	<p><b>Cummings, J. G., &amp; Wittenberg, J. V.</b> (2008). Supportive expressive therapy - Parent child version: an exploratory study. <i>Psychotherapy Theory, Research, Practice, Training</i>, 45, 148-164. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22122414">http://www.ncbi.nlm.nih.gov/pubmed/22122414</a></p>	<b>Oppositional defiant disorder/disruptive behavior disorder</b> (children)	<b>Supportive Expressive Therapy – Parent Child</b> (SET-PC; Wittenberg, adapted from Luborsky, 19854)	<b>The Incredible Years Parenting Program</b>
50.	<p><b>Dare, C.,</b> Eisler, I., Russell, G., Treasure, J., &amp; Dodge, L. (2001). Psychological therapies for adults with anorexia nervosa: randomised controlled trial of out-patient treatments. <i>Br J Psychiatry</i>, 178, 216-221. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11230031">http://www.ncbi.nlm.nih.gov/pubmed/11230031</a></p>	<b>Anorexia</b>	<b>Focal psychoanalytic psychotherapy</b> (Dare, 1995; Malan, 1979)	<p><b>Cognitive Analytic Therapy</b> (Ryle, 1990; Treasure et al, 1995)</p> <p><b>Family therapy</b> (no specific reference)</p>
51.	<p><b>de Jonghe, F.,</b> Hendriksen, M., van Aalst, G., Kool, S., Peen, J., &amp; Van, R., et al. (2004). Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. <i>British Journal of Psychiatry</i>, 185, 37-45.</p>	<b>Depression</b>	<b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, 2005)	<b>Short Psychodynamic Supportive Psychotherapy + antidepressants</b>

	<a href="http://www.ncbi.nlm.nih.gov/pubmed/15231554">http://www.ncbi.nlm.nih.gov/pubmed/15231554</a>			
52.	<b>de Jonghe, F., Kool, S., van Aalst, G., Dekker, J., &amp; Peen, J. (2001).</b> Combining psychotherapy and antidepressants in the treatment of depression. <i>Journal of Affective Disorders, 64</i> (2-3), 217–29. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11313088">http://www.ncbi.nlm.nih.gov/pubmed/11313088</a>	<b>Depression</b>	<b>Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) + medication</b>	<b>Medication alone</b>
53.	<b>de Roten, Y., Ambresin, G., Herrera, F., Fassassi, S., Fournier, N., Preisig, M., &amp; Despland, J.-N. (2017).</b> Efficacy of an adjunctive brief psychodynamic psychotherapy to usual inpatient treatment of depression: Results of a randomized controlled trial. <i>Journal of Affective Disorders, 209</i> , 105–113. <a href="http://doi.org/10.1016/j.jad.2016.11.013">http://doi.org/10.1016/j.jad.2016.11.013</a>	<b>Depression</b>	<b>Intensive Brief Psychodynamic Psychotherapy (IBPP; Despland et al., 2010) + TAU</b>	<b>TAU only</b>
54.	<b>Dekker, J. J. M., Koelen, J. A., Van, H. L., Schoevers, R. A., Peen, J., Hendriksen, M., ... De Jonghe, F. (2008).</b> Speed of action: the relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. <i>Journal of Affective Disorders, 109</i> (1-2), 183–188. <a href="http://linkinghub.elsevier.com/retrieve/pii/S0165032707003588">http://linkinghub.elsevier.com/retrieve/pii/S0165032707003588</a>  <b>Dekker, J., Van, H. L., Hendriksen, M., Koelen, J., Schoevers, R. A., Kool, S., ... Peen, J. (2013).</b> What is the best sequential treatment strategy in the treatment of depression? Adding pharmacotherapy to psychotherapy or vice versa? <i>Psychotherapy and Psychosomatics, 82</i> (2), 89–98. <a href="http://www.karger.com/doi/10.1159/000341177">http://www.karger.com/doi/10.1159/000341177</a>	<b>Depression</b>	<b>Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005)</b>	<b>Pharmacotherapy</b>
55.	<b>Dekker, J., Molenaar, P. J., Kool, S., Van Aalst, G., Peen, J., &amp; de Jonghe, F. (2005).</b> Dose–effect relations in time-limited combined psycho-pharmacological treatment for depression. <i>Psychological Medicine, 35</i> (1), 47–58. <a href="http://www.journals.cambridge.org/abstract/S0033291704002685">http://www.journals.cambridge.org/abstract/S0033291704002685</a>	<b>Depression</b>	<b>Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) 8 sessions + medication</b>	<b>Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) 16 sessions + medication</b>
56.	<b>Dinger, U., Klipsch, O., Köhling, J., Ehrenthal, J. C., Nikendei, C., Herzog, W., &amp; Schauenburg, H. (2014).</b> Day-clinic and inpatient psychotherapy for depression (DIP-D): A randomized controlled pilot study in routine clinical care. <i>Psychotherapy and Psychosomatics, 83</i> (3), 194–195. <a href="http://doi.org/10.1159/000357437">http://doi.org/10.1159/000357437</a>	<b>Depression</b>	<b>Day-clinic multimodal psychodynamic psychotherapy (no specific manual referenced)</b>	<b>Inpatient multimodal psychodynamic psychotherapy (no specific manual referenced)</b>
57.	<b>Doering, S., Hörz, S., Rentrop, M., Fischer-Kern, M., Schuster, P., &amp; Benecke, C., et al. (2010).</b> Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: randomised controlled trial. <i>Br J Psychiatry, 196</i> , 389–395. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20435966">http://www.ncbi.nlm.nih.gov/pubmed/20435966</a>	<b>Borderline personality disorder</b>	<b>Transference-focused psychotherapy (TFP; Kernberg)</b>	<b>TAU (experienced therapists)</b>
58.	<b>Driessen, E., Van, L. H., Don, F. J., Peen, J.,</b>	<b>Depression</b>	<b>Short-term</b>	<b>Cognitive-Behavior</b>

	<p>Kool, S., Westra, D., ... Dekker, J. J. M. (2013). The Efficacy of Cognitive-Behavioral Therapy and Psychodynamic Therapy in the Outpatient Treatment of Major Depression: A Randomized Clinical Trial. <i>American Journal of Psychiatry</i>, 170(9), 1041-1050 <a href="http://ajp.psychiatryonline.org/doi/abs/10.1176/appi.ajp.2013.12070899">http://ajp.psychiatryonline.org/doi/abs/10.1176/appi.ajp.2013.12070899</a></p> <p>Driessen, E., Van, H. L., Peen, J., Don, F. J., Kool, S., Westra, D., ... Dekker, J. J. (2014). Therapist-rated outcomes in a randomized clinical trial comparing cognitive behavioral therapy and psychodynamic therapy for major depression. <i>Journal of Affective Disorders</i>, 170C, 112–118. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25240140">http://www.ncbi.nlm.nih.gov/pubmed/25240140</a></p> <p>Driessen, E., Van, H. L., Peen, J., Don, F. J., Twisk, J. W. R., Cuijpers, P., &amp; Dekker, J. J. M. (2017). Cognitive-Behavioral Versus Psychodynamic Therapy for Major Depression: Secondary Outcomes of a Randomized Clinical Trial. <i>Journal of Consulting &amp; Clinical Psychology</i>, 85(7), 653–663. <a href="http://dx.doi.org/10.1037/ccp0000207">http://dx.doi.org/10.1037/ccp0000207</a></p>		<p><b>psychodynamic supportive psychotherapy</b> (de Jonghe, 2005)</p>	<p><b>Therapy</b> (CBT; Molenaar et al., 2009)</p>
59.	<p><b>Durham</b>, R. C., Guthrie, M., Morton, R. V., Reid, D. a, Treliving, L. R., Owler, D. F., &amp; Macdonald, R. A. R. (2003). Tayside-Fife clinical trial of cognitive-behavioural therapy for medication-resistant psychotic symptoms. <i>British Journal of Psychiatry</i>, 182, 303–311. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12668405">http://www.ncbi.nlm.nih.gov/pubmed/12668405</a></p>	<p><b>Schizophrenia</b> (and other medication-resistant psychotic symptoms)</p>	<p><b>Supportive therapy</b> (dynamically oriented; Garfield, 1995; De Jonghe, 1993)</p>	<p><b>CBT</b> (Tarrier, 1992; Kingdon &amp; Turkington, 1994). <b>+ TAU</b></p>
60.	<p><b>Durham</b>, R. C., Murphy, T., Allan, T., Richard, K., Treliving, L. R., &amp; Fenton, G. W., et al. (1994). Cognitive therapy, analytic psychotherapy and anxiety management training for generalised anxiety disorder. <i>Br J Psychiatry</i>, 165, 315-323. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7994500">http://www.ncbi.nlm.nih.gov/pubmed/7994500</a></p>	<p><b>Generalized anxiety disorder (GAD)</b></p>	<p><b>Analytic-based psychotherapy</b> (no specific reference)</p>	<p><b>Cognitive Therapy / Anxiety Management training</b> (Beck &amp; Emery, 1985)</p>
61.	<p><b>Edginton</b>, E., Walwyn, R., Twiddy, M., Wright-hughes, A., Tubeuf, S., Reed, S., ... Group, F. (2018). TIGA-CUB-manualised psychoanalytic child psychotherapy versus treatment as usual for children aged 5 – 11 with treatment-resistant conduct disorders and their primary carers: results from a randomised controlled feasibility trial. <i>Journal of Child and Adolescent Mental Health</i>, 0583. <a href="http://doi.org/10.2989/17280583.2018.1532433">http://doi.org/10.2989/17280583.2018.1532433</a></p>	<p><b>Treatment-resistant conduct disorders</b> (children and carers)</p>	<p><b>Manualised psychoanalytic child psychotherapy</b> (mPCP)</p>	<p><b>TAU</b></p>
62.	<p><b>Emmelkamp</b>, P. M. G., Benner, A., Kuipers, A., Feiertag, G. A., Koster, H. C., &amp; Van Apeldoorn, F. J., et al. (2006). Comparison of brief dynamic and cognitive-behavioural therapies in avoidant personality disorder. <i>Br J Psychiatry</i>, 189, 60-64. <a href="http://www.ncbi.nlm.nih.gov/pubmed/16816307">http://www.ncbi.nlm.nih.gov/pubmed/16816307</a></p>	<p><b>Avoidant personality disorder</b></p>	<p><b>Brief Dynamic Therapy</b> (BDT; Malan, 1979; also reference to Luborsky, 1984)</p>	<p><b>CBT</b> (Beck &amp; Freeman, 1990; Emmelkamp et al, 1992)</p>
63.	<p><b>Fairburn</b>, G., Kirk, J., Connor, M. O., &amp; Cooper, P. J. (1986). A comparison of two psychological treatments for bulimia. <i>Behaviour research and therapy</i>, 24(6), 629–643.</p>	<p><b>Bulimia</b></p>	<p><b>Short-term Focal Psychotherapy</b> (Rosen, 1979)</p>	<p><b>CBT</b> (Fairburn, 1985)</p>

	<a href="http://www.sciencedirect.com/science/article/pii/S0005796786900586">http://www.sciencedirect.com/science/article/pii/S0005796786900586</a>			
64.	<p><b>Faramarzi, M., Azadfallah, P., Book, H. E., Tabatabaei, K. R., Taheri, H., &amp; Shokri-Shirvani, J. (2013).</b> A randomized controlled trial of brief psychoanalytic psychotherapy in patients with functional dyspepsia. <i>Asian journal of psychiatry</i>, 6(3), 228–34. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23642981">http://www.ncbi.nlm.nih.gov/pubmed/23642981</a></p> <p>Faramarzi, M., &amp; Book, H. E. (2015). The Effect of Psychotherapy in Improving Physical and Psychiatric Symptoms in Patients with Functional Dyspepsia, <i>Iran Journal of Psychiatry</i>, 43–49. <a href="http://www.ncbi.nlm.nih.gov/pubmed/26005480">http://www.ncbi.nlm.nih.gov/pubmed/26005480</a></p>	<b>Functional dyspepsia</b>	<b>Brief psychodynamic therapy</b> (Book, 1998) based on Luborsky (1984) + standard medical therapy	<b>TAU</b> (medical therapy)
65.	<p><b>Fonagy, P., Rost, F., Carlyle, J.-A., Mchpherson, S., Thomas, R., Fearon, R. M. P., ... Taylor, D. (2015).</b> Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression : the Tavistock Adult Depression Study (TADS ). <i>World Psychiatry</i>, 14(3), 312-321 <a href="http://www.ncbi.nlm.nih.gov/pubmed/26407787">http://www.ncbi.nlm.nih.gov/pubmed/26407787</a></p>	<b>Depression</b> (treatment resistant)	<b>Long-Term Psychoanalytic Psychotehrapy</b> (LTPP; Taylor, 2015)	<b>TAU</b>
66.	<p><b>Fonagy, P., Slead, M., &amp; Baradon, T. (2016).</b> Randomized Controlled Trial of Parent-Infant Psychotherapy for Parents With Mental Health Problems and Young Infants. <i>Infant Mental Health Journal</i>, 37(2), 97-114 <a href="http://doi.wiley.com/10.1002/imhj.21553">http://doi.wiley.com/10.1002/imhj.21553</a></p>	<b>Parents with mental health problems</b>	<b>Parent-infant psychotherapy (PIP)</b>	<b>Standard care</b>
67.	<p><b>Gallagher-Thompson, D., &amp; Steffen, A. M. (1994).</b> Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers. <i>J Consult Clin Psychol</i>, 62, 543-549. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8063980">http://www.ncbi.nlm.nih.gov/pubmed/8063980</a></p>	<b>Depression</b> (among family caregivers)	<b>Brief Psychodynamic Therapy</b> (Mann, 1973)	<b>CBT</b> (Beck et al., 1979; Lewinsohn et al, 1985; Gallagher et alm 1981)
68.	<p><b>Gallagher, D. E., &amp; Thompson, L. W. (1982).</b> Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. <i>Psychotherapy: Theory, Research, and Practice</i>, 482-490. <a href="http://psycnet.apa.org/journals/pst/19/4/482">http://psycnet.apa.org/journals/pst/19/4/482</a></p>	<b>Depression</b>	<b>Brief relational/insight psychotherapy</b> (Bellak & Small, 1965)	<b>Behavioral therapy / Cognitive therapy</b> (Lewinsohn et al, 1976; Gallagher et al., 1981)
69.	<p><b>Garner, D. M., Rockert, W., Davis, R., Garner, M. V., Olmsted, M. P., &amp; Eagle, M., et al. (1993).</b> Comparison of cognitive-behavioral and supportive-expressive therapy for bulimia nervosa. <i>Am J Psychiatry</i>, 150, 37-46. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8417578">http://www.ncbi.nlm.nih.gov/pubmed/8417578</a></p>	<b>Bulimia</b>	<b>Supportive-expressive psychotherapy</b> (Luborsky, 1984)	<b>CBT</b> (Fairburn, 1985)
70.	<p><b>Gelder, M. G., Marks, I. M., &amp; Wolff, H. H. (1967).</b> Desensitization and psychotherapy in the treatment of phobic states: a controlled inquiry. <i>British Journal of Psychiatry</i>, 113(494), 53–81. <a href="http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.113.494.53">http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.113.494.53</a></p>	<b>Phobic states</b>	<b>Psychotherapy</b> (no reference)	<b>1. Desensitization</b> (Wolpe)  <b>2. Group psychotherapy</b> (Foulkes)
71.	<p><b>Gersons, B. P., Carlier, I. V, Lamberts, R. D., &amp; van der Kolk, B. a. (2000).</b> Randomized clinical</p>	<b>PTSD</b>	<b>Brief eclectic psychotherapy</b> (BEP;	<b>Waitlist</b>

	<p>trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder. <i>Journal of Traumatic Stress</i>, 13(2), 333–47. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10838679">http://www.ncbi.nlm.nih.gov/pubmed/10838679</a></p>		Gersons & Carlier, 1994; Manual includes both CBT and PDT components)	
72.	<p><b>Ghorbani, N., Dadsetan, P., Ejei, J., &amp; Motiyan, H.</b> (2000). The consequences of overcoming resistance and emotional disclosure on lymphocyte T-helper and T-suppressor and psychological pathology. <i>Journal of Psychology (in persian)</i>, 3, 368–89.</p>	<b>Students</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Davanloo, 1990)	<b>Simple verbal disclosure groups (SVD)</b>
73.	<p><b>Giesen-Bloo, J., van Dyck, R., Spinhoven, P., van Tilburg, W., Dirksen, C., &amp; van Asselt, T., et al.</b> (2006). Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. <i>Arch Gen Psychiatry</i>, 63, 649-658. <a href="http://www.ncbi.nlm.nih.gov/pubmed/16754838">http://www.ncbi.nlm.nih.gov/pubmed/16754838</a></p> <p>van Asselt, A. D. I., Dirksen, C. D., Arntz, A., Giesen-Bloo, J. H., van Dyck, R., Spinhoven, P., ... Severens, J. L. (2008). Out-patient psychotherapy for borderline personality disorder: cost-effectiveness of schema-focused therapy v. transference-focused psychotherapy. <i>The British Journal of Psychiatry: The Journal of Mental Science</i>, 192(6), 450–7. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18515897">http://www.ncbi.nlm.nih.gov/pubmed/18515897</a></p>	<b>Borderline personality disorder</b>	<b>Transference focused psychotherapy</b> (TFP; Clarkin, Yeomans, & Kernberg, 1999; Yeomans, Clarkin, & Kernberg, 2002)	<b>Schema-focused therapy</b> (Young, 1994; Young et al, 2003)
74.	<p><b>Gilboa-Schechtman, E., Foa, E. B., Shafraan, N., Aderka, I. M., Powers, M. B., Rachamim, L., ... Apter, A.</b> (2010). Prolonged exposure versus dynamic therapy for adolescent PTSD: a pilot randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i>, 49(10), 1034–42. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2945239&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2945239&amp;tool=pmcentrez&amp;rendertype=abstract</a></p>	<b>PTSD (adolescents)</b>	<b>Time-limited dynamic therapy</b> (TLDP-A; Shefner et al, 1995)	<b>Prolonged exposure (Foa)</b>
75.	<p><b>Goodyer, I. M., Reynolds, S., Barrett, B., Byford, S., Dubicka, B., Hill, J., ... Midgley, N.</b> (2016). Cognitive behavioural therapy and short-term psychoanalytical psychotherapy versus a brief psychosocial intervention in adolescents with unipolar major depressive disorder (IMPACT): a multicentre, pragmatic, observer-blind, randomised controlled superiority trial. <i>The Lancet Psychiatry</i>, 366(16), 1–11. <a href="http://doi.org/10.1016/S2215-0366(16)30378-9">http://doi.org/10.1016/S2215-0366(16)30378-9</a></p>	<b>Depression (adolescents)</b>	<b>Short-Term Psychoanalytic Psychotherapy</b> (STPP; Cregeen et al., 2016)	<p><b>1. CBT</b> (Graham et al., 2015)</p> <p><b>2. Brief Psychosocial Intervention</b> (BPI; Trowell et al., 2009)</p>
76.	<p><b>Gowers, S., Norton, K., Halek, C., &amp; Crisp, A. H.</b> (1994). Outcome of outpatient psychotherapy in a random allocation treatment study of anorexia nervosa. <i>Int J Eat Disord</i>, 15, 165-177. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8173562">http://www.ncbi.nlm.nih.gov/pubmed/8173562</a></p>	<b>Anorexia (adolescents and young women)</b>	<b>Short term individual + family interviews</b> (Koss & Butcher, 1986)	<b>Assessment only</b>
77.	<p><b>Gregory, R. J., Chlebowski, S., Kang, D., Remen, A. L., Soderberg, M. G., &amp; Stepkovitch, J., et al.</b> (2008). A controlled trial of psychodynamic psychotherapy for co-</p>	<b>Borderline personality disorder + alcohol use disorder</b>	<b>Dynamic deconstructive therapy</b> (Gregory & Remen)	<b>TAU</b>

	occurring borderline personality disorder and alcohol use disorder. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 45, 28-41. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.45.1.28">http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.45.1.28</a>			
78.	<b>Grothe, C., Bering, R., SpieB, R., Luneburg, E., &amp; Fischer, G. (2003).</b> Mehrdimensionale Psychodynamische Traumaterapie (MPTT) Forschungsergebnisse zur Standardversion. <i>Zeitschrift für Psychotraumatologie, Psychotherapiewissenschaft und Psychologische Medizin</i> 1(2), 27–43.	<b>PTSD</b>	<b>Mehrdimensionale Psychodynamische Traumaterapie (MPTT;</b> Fischer, 2000)	<b>1. EMDR (Shapiro, 1995; 1998)</b> <b>2. Waitlist control (?)</b>
79.	<b>Gunderson, J. G., Frank, A. F., Katz, H. M., Vannicelli, M. L., Frosch, J. P., &amp; Knapp, P. H., et al. (1984).</b> Effects of psychotherapy in schizophrenia: II. Comparative outcome of two forms of treatment. <i>Schizophr Bull</i> , 10, 564-598. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6151246">http://www.ncbi.nlm.nih.gov/pubmed/6151246</a>	<b>Schizophrenia</b>	<b>Exploratory, insight oriented psychotherapy</b> (Adapted from Fromm-Reichmann)	<b>Reality-adaptive, supportive psychotherapy</b>
80.	<b>Guthrie, E., Creed, F., Dawson, D., &amp; Tomenson, B. (1991).</b> A controlled trial of psychological treatment for the irritable bowel syndrome. <i>Gastroenterology</i> , 100, 450-457. <a href="http://www.gastrojournal.org/article/0016-5085(91)90215-7/pdf">http://www.gastrojournal.org/article/0016-5085(91)90215-7/pdf</a>  <b>Guthrie, E., Creed, F., Dawson, D., &amp; Tomenson, B. (1993).</b> A randomised controlled trial of psychotherapy in patients with refractory Irritable Bowel Syndrome. <i>British Journal of Psychiatry</i> , 163, 315–21. <a href="http://bjp.rcpsych.org/content/163/3/315">http://bjp.rcpsych.org/content/163/3/315</a>	<b>Irritable bowel syndrome</b>	<b>Psychodynamic-Interpersonal Therapy (PIT;</b> Hobson, 1985)	<b>Supportive listening</b>
81.	<b>Guthrie, E., Kapur, N., Mackway-Jones, K., Chew-Graham, C., Moorey, J., &amp; Mendel, E., et al. (2001).</b> Randomised controlled trial of brief psychological intervention after deliberate self poisoning. <i>BMJ</i> , 323, 135-138. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=34723&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=34723&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Self-poisoning</b>	<b>Psychodynamic-Interpersonal Therapy (PIT;</b> Hobson, 1985)	<b>TAU</b>
82.	<b>Guthrie, E., Moorey, J., Margison, F., Barker, H., Palmer, S., &amp; McGrath, G., et al. (1999).</b> Cost-effectiveness of brief psychodynamic-interpersonal therapy in high utilizers of psychiatric services. <i>Arch Gen Psychiatry</i> , 56, 519-526. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10359466">http://www.ncbi.nlm.nih.gov/pubmed/10359466</a>	<b>Mixed sample</b>	<b>Psychodynamic-Interpersonal Therapy (PIT;</b> Hobson, 1985)	<b>TAU</b>
83.	<b>Hall, A., &amp; Crisp, A. H. (1987).</b> Brief psychotherapy in the treatment of anorexia nervosa: Outcome at one year. <i>British Journal of Psychiatry</i> , 151, 185-191. <a href="http://www.ncbi.nlm.nih.gov/pubmed/3690108">http://www.ncbi.nlm.nih.gov/pubmed/3690108</a>	<b>Anorexia</b>	<b>Combined individual and family psychotherapy (no reference)</b>	<b>Dietary advice</b>
84.	<b>Hamilton, J., Guthrie, E., Creed, F., Thompson, D., Tomenson, B., &amp; Bennett, R., et al. (2000).</b> A randomized controlled trial of psychotherapy in patients with chronic functional dyspepsia. <i>Gastroenterology</i> , 119, 661-669. <a href="http://linkinghub.elsevier.com/retrieve/pii/S0016508500145949">http://linkinghub.elsevier.com/retrieve/pii/S0016508500145949</a>	<b>Chronic Functional Dyspepsia</b>	<b>Psychodynamic-Interpersonal Therapy (PIT;</b> Hobson, 1985)	<b>Supportive therapy (Guthrie, unpub.)</b>

85.	<p><b>Hardy, G. E., Barkham, M., Shapiro, D. a, Stiles, W. B., Rees, a, &amp; Reynolds, S. (1995).</b> Impact of Cluster C personality disorders on outcomes of contrasting brief psychotherapies for depression. <i>Journal of consulting and clinical psychology, 63(6)</i>, 997–1004.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/8543722">http://www.ncbi.nlm.nih.gov/pubmed/8543722</a></p>	<p><b>Depression + personality disorder cluster C</b></p>	<p><b>Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)</b></p>	<p><b>CBT (Firth &amp; Shapiro, 1985)</b></p>
86.	<p><b>Hellerstein, D. J., Rosenthal, R. N., Pinsker, H., Samstag, L. W., Muran, J. C., &amp; Winston, A., et al. (1998).</b> A randomized prospective study comparing supportive and dynamic therapies. Outcome and alliance. <i>J Psychother Pract Res, 7</i>, 261-271.  <a href="http://psycnet.apa.org/psycinfo/1998-12001-001">http://psycnet.apa.org/psycinfo/1998-12001-001</a></p>	<p><b>Personality disorders (cluster C)</b></p>	<p><b>Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1980)</b></p>	<p><b>Brief Adaptive Therapy (Winston?)</b></p>
87.	<p><b>Herrmann-Lingen, C., Beutel, M. E., Bosbach, A., Deter, H., Fritzsche, K., Hellmich, M., ... Albus, C. (2016).</b> Stepwise Psychotherapy Intervention for Reducing Risk in Coronary Artery Disease: Results of an Observer-Blinded, Multicenter, Randomized Trial in Depressed Patients With Coronary Artery Disease. <i>Psychosomatic Medicine, 1–12</i>.  <a href="http://doi.org/10.1097/PSY.0000000000000332">http://doi.org/10.1097/PSY.0000000000000332</a></p>	<p><b>Depressive symptoms in patients with coronary artery disease</b></p>	<p><b>Stepwise psychotherapy program (3 sessions supportive-expressive psychotherapy (Luborsky, 1984) + group therapy (no specific manual mentioned)</b></p>	<p><b>TAU</b></p>
88.	<p><b>Hersen, M., Himmelhoch, J. M., Thase, M. E., &amp; Bellack, A. S. (1984).</b> Effects of social skill training, amitriptyline, and psychotherapy in unipolar depressed women. <i>Behavior Therapy, 15(1)</i>, 21–40.  <a href="http://linkinghub.elsevier.com/retrieve/pii/S005789484800398">http://linkinghub.elsevier.com/retrieve/pii/S005789484800398</a></p>	<p><b>Depression</b></p>	<p><b>Time-limited dynamic therapy (no reference) + placebo</b></p>	<p><b>1. Social skill training + placebo</b>  <b>2. Amitriptyline</b></p>
89.	<p><b>Hertzmman, L., Target, M., Hewison, D., Casey, P., Fearon, P., &amp; Lassri, D. (2016).</b> Mentalization-based therapy for parents in entrenched conflict: A random allocation feasibility study. <i>Psychotherapy, 53(4)</i>, 388-401. <a href="http://psycnet.apa.org/buy/2016-58498-002">http://psycnet.apa.org/buy/2016-58498-002</a></p>	<p><b>Parents in entrenched conflict</b></p>	<p><b>Mentalization-Based Treatment for Parental Conflict (MBT-PT; Hertzmman &amp; Abse, 2008; Nyberg &amp; Hertzmman, 2014)</b></p>	<p><b>Parent Group (psychoeducation)</b></p>
90.	<p><b>Hoffart, A., &amp; Martinsen, E. W. (1990).</b> Exposure-based integrated vs. Pure psychodynamic treatment of agoraphobic inpatients. <i>Psychotherapy: Theory, Research, Practice, Training, Vol 27(2)</i>, 210–218.  <a href="http://psycnet.apa.org/psycinfo/1990-29285-001">http://psycnet.apa.org/psycinfo/1990-29285-001</a></p>	<p><b>Agoraphobia</b></p>	<p><b>Integrative behavioural-psychodynamic treatment</b></p>	<p><b>Psychodynamic oriented group therapy</b></p>
91.	<p><b>Høglend, P., Amlo, S., Marble, A., Bøggwald, K. P., Sørbye, O., &amp; Sjaastad, M. C., et al. (2006).</b> Analysis of the patient-therapist relationship in dynamic psychotherapy: an experimental study of transference interpretations. <i>Am J Psychiatry, 163</i>, 1739-1746.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/17012684">http://www.ncbi.nlm.nih.gov/pubmed/17012684</a></p> <p>Høglend, P., Bøggwald, K.-P., Amlo, S., Marble, A., Ulberg, R., Sjaastad, M. C., ... Johansson, P. (2008). Transference interpretations in dynamic psychotherapy: do they really yield sustained effects? <i>The American Journal of Psychiatry, 165(6)</i>, 763–71.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/18413707">http://www.ncbi.nlm.nih.gov/pubmed/18413707</a></p>	<p><b>Mixed sample</b></p>	<p><b>Psychodynamic psychotherapy with transference interpretation (Høglend, 1995)</b></p>	<p><b>Psychodynamic psychotherapy without transference interpretation (Høglend, 1995)</b></p>

92.	<p><b>Horn, H., Geiser-Elze, A., Reck, C., Hartmann, M., Victor, D., Winkelmann, K., &amp; Kronmüller, K.-T. (2005).</b> Zur Wirksamkeit psychodynamischer Kurzzeitpsychotherapie bei Kindern und Jugendlichen mit Angststörungen [Efficacy of psychodynamic short-term psychotherapy for children and adolescents with depression]. <i>Praxis Der Kinderpsychologie Und Kinderpsychiatrie</i>, 54, 578–597. Article in German. Retrieved from <a href="http://search.ebscohost.com/login.aspx?direct=true&amp;db=psyh&amp;AN=2005-13295-002&amp;site=ehost-live">http://search.ebscohost.com/login.aspx?direct=true&amp;db=psyh&amp;AN=2005-13295-002&amp;site=ehost-live</a></p>	<b>Depression</b>	<b>Psychodynamic Short-Term Psychotherapy</b> (PSTP: unpublished manual by Hartmann, Horn, & Kronmüller)	<b>Waitlist</b>
93.	<p><b>Huber, D., Henrich, G., Gastner, J., &amp; Klug, G. (2012).</b> Must All Have Prizes? The Munich Psychotherapy Study. In <i>Psychodynamic Psychotherapy Research</i>. <a href="http://link.springer.com/chapter/10.1007%2F978-1-60761-792-1_4#page-1">http://link.springer.com/chapter/10.1007%2F978-1-60761-792-1_4#page-1</a></p> <p>Huber, D., Huber, D., Zimmermann, J., Henrich, G., &amp; Klug, G. (2012). Comparison of cognitive-behaviour therapy with psychoanalytic and psychodynamic therapy for depressed patients – A three-year follow-up study. <i>Zeitschrift für Psychosomatic</i>, (58), 299–316. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22987495">http://www.ncbi.nlm.nih.gov/pubmed/22987495</a></p> <p>Huber, D., Henrich, G., Clarkin, J., &amp; Klug, G. (2013). Psychoanalytic versus psychodynamic therapy for depression: a three-year follow-up study. <i>Psychiatry</i>, 76(2), 132–49. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23631544">http://www.ncbi.nlm.nih.gov/pubmed/23631544</a></p>	<b>Depression</b>	<b>Psychoanalysis</b> (non-manualized)	<p><b>1. LTPP</b> (non-manualized)</p> <p><b>2. CBT</b> (non-manualized; patients not randomly assigned to this group)</p>
94.	<p><b>Hubschmid, M., Aybek, S., D, M., Maccaferri, G. E., D, M., Chocron, O., ... D, M. (n.d.).</b> Efficacy of brief interdisciplinary psychotherapeutic intervention for motor conversion disorder and nonepileptic attacks. <i>General Hospital Psychiatry</i>, 37(5), 448–455. doi:10.1016/j.genhosppsych.2015.05.007 <a href="http://dx.doi.org/10.1016/j.genhosppsych.2015.05.007">http://dx.doi.org/10.1016/j.genhosppsych.2015.05.007</a></p>	<b>Motor conversion disorder and nonepileptic seizures</b>	<b>Interdisciplinary psychotherapeutic intervention</b> (Psychodynamic-Interpersonal therapy based on Guthrie, 1999)	<b>Standard care</b>
95.	<p><b>Jakobsen, J. C., Gluud, C., Kongerslev, M., Larsen, K. A., Sørensen, P., Winkel, P., ... Simonsen, E. (2014).</b> Third-wave cognitive therapy versus mentalisation-based treatment for major depressive disorder: A randomised clinical trial. <i>BMJ Open</i>, 4(8), 1–12. <a href="http://doi.org/10.1136/bmjopen-2014-004903">http://doi.org/10.1136/bmjopen-2014-004903</a></p>	<b>Depression</b>	<b>Mentalization-Based Treatment</b> (MBT; Larsen KA. Mentalisation-based therapy. A treatment manual (in Danish). 2009. <a href="http://www.ctu.dk">http://www.ctu.dk</a> )	<b>Third Wave Cognitive Therapy</b> ( <a href="http://ctu.dk/publications/supplementary-material.aspx">http://ctu.dk/publications/supplementary-material.aspx</a> ).
96.	<p><b>Jantschek, G., Zeitz, M., Wirsching, M., Klor, H. U., Studt, H. H., Rasenack, J., ... Disease, T. G. S. G. on P. I. in C. (1999).</b> Effect of psychotherapy on the course of Crohn's disease. <i>Psychosomatic Intervention in CD</i>, 1289–1296. <a href="http://www.ncbi.nlm.nih.gov/pubmed/9930393">http://www.ncbi.nlm.nih.gov/pubmed/9930393</a></p> <p>Keller, W., Pritsch, M., Von Wietersheim, J., Scheib, P., Osborn, W., &amp; Balck, F., et al. (2004). Effect of psychotherapy and relaxation on the psychosocial and somatic course of Crohn's disease: main results of the German</p>	<b>Crohn's disease</b>	<b>PDT+relaxation+drug treatment</b> (no manual was used but reference to Luborsky, 1984)	<b>Drug treatment</b>



	Prospective Multicenter Psychotherapy Treatment study on Crohn's Disease. <i>J Psychosom Res</i> , 56, 687-696. <a href="http://www.ncbi.nlm.nih.gov/pubmed/15193965">http://www.ncbi.nlm.nih.gov/pubmed/15193965</a>			
97.	<b>Johansson, R., Björklund, M., Hornborg, C., Karlsson, S., Hesser, H., Ljótsson, B., ... Andersson, G. (2013).</b> Affect-focused psychodynamic psychotherapy for depression and anxiety through the Internet: a randomized controlled trial. <i>PeerJ</i> , 1, e102. <a href="https://peerj.com/articles/102">https://peerj.com/articles/102</a>	<b>Depression and/or anxiety</b>	<b>IPDT</b> (Affect-focused; based on Frederick, 2009)	<b>Online therapist support and clinical monitoring</b>
98.	<b>Johansson, R., Ekbladh, S., Hebert, A., Lindström, M., Möller, S., Pettit, E., ... Andersson, G. (2012).</b> Psychodynamic Guided Self-Help for Adult Depression through the Internet: A Randomised Controlled Trial. (A. Bruce, Ed.) <i>PLoS ONE</i> , 7(5), <a href="http://dx.plos.org/10.1371/journal.pone.0038021">http://dx.plos.org/10.1371/journal.pone.0038021</a>	<b>Depression</b>	<b>IPDT</b> (SUBGAP; based on Silverberg, 2005)	<b>Psychoeducation + support</b>
99.	<b>Johansson, R., Hesslow, T., Ljótsson, B., Jansson, A., Jonsson, L., Färdig, S., Karlsson, J., Hesser, H., Frederik, R. J., Lilliengren, P., Carlbring, P., &amp; Andersson, G. (2017).</b> Internet-based affect-focused psychodynamic therapy for social anxiety disorder: A randomized controlled trial with 2-year follow-up. <i>Psychotherapy</i> , 54(4). <a href="http://doi.org/10.1037/pst0000147">http://doi.org/10.1037/pst0000147</a>	<b>Social anxiety disorder (SAD)</b>	<b>IPDT</b> (Affect-focused; based on Frederick, 2009)	<b>Waitlist</b>
100.	<b>Jørgensen, C. R., Freund, C., Bøye, R., Jordet, H., Andersen, D., &amp; Kjølbye, M. (2012).</b> Outcome of mentalization-based and supportive psychotherapy in patients with borderline personality disorder: a randomized trial. <i>Acta psychiatrica Scandinavica</i> , 1–13. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22897123">http://www.ncbi.nlm.nih.gov/pubmed/22897123</a>	<b>Borderline personality disorder</b>	<b>Mentalization-Based Therapy</b> (MBT; Bateman & Fonagy, 2001)	<b>Supportive group therapy</b>
101.	<b>Kamkhagi, D., Costa, A. C. O., Kusminsky, S., Supino, D., Diniz, B. S., Gattaz, W. F., &amp; Forlenza, O. V. (2015).</b> Benefits of psychodynamic group therapy on depression, burden and quality of life of family caregivers to Alzheimer's disease patients. <i>Archives of Clinical Psychiatry</i> , 42(6), 157–160. <a href="http://doi.org/10.1016/j.jalz.2014.05.934">http://doi.org/10.1016/j.jalz.2014.05.934</a>	<b>Family caregivers to Alzheimer's disease</b> (depression, quality of life, burden)	<b>Psychodynamic Group Psychotherapy</b> (PGT)	<b>Body Awareness Therapy (BAT)</b>
102.	<b>Karon, B. P., &amp; Vandenbos, G. R. (1972).</b> The consequences of psychotherapy for schizophrenic patients. <i>Psychotherapy: Theory, Research &amp; Practice</i> , 9(2), 111–119. <a href="http://doi.org/10.1037/h0086728">http://doi.org/10.1037/h0086728</a>	<b>Schizophrenia</b>	<b>Psychoanalytic psychotherapy</b> (Karon, 1963, 1964; Karon & Rosberg, 1958; Rosberg & Karon, 1958)	<b>1. Psychoanalytic psychotherapy</b> ("Ego-oriented"; Tierney, 1965) + <b>medication</b> <b>2. Medication alone</b>
103.	<b>Klein, D. F., Zitrin, C. M., Woerner, M. G., &amp; Ross, D. C. (1983).</b> Treatment of Phobias. II. Behavior therapy and supportive psychotherapy: are there any specific ingredients? <i>Archives of General Psychiatry</i> , 139–145. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6130751">http://www.ncbi.nlm.nih.gov/pubmed/6130751</a>	<b>Phobias</b>	<b>Supportive therapy</b> (dynamically oriented; no reference)	<b>Behavior therapy</b> (no reference)
104.	<b>Kleinman, P. H., Woody, G. E., Todd, T. C., Millman, R. B., Kang, S. Y., &amp; Kemp, J., et al. (1990).</b> Crack and cocaine abusers in outpatient psychotherapy. <i>NIDA Res Monogr</i> ,	<b>Cocaine abuse</b>	<b>Supportive-Expressive Therapy</b> (Luborsky, 1984)	<b>Family therapy + para professionally led group therapy</b>

	104, 24-35.			
105.	<p><b>Knekt, P., &amp; Lindfors, O. (2004).</b> A randomized trial of the effect of four forms of psychotherapy on depressive and anxiety disorders. Helsinki: Kela. <a href="https://helda.helsinki.fi/handle/10250/3315">https://helda.helsinki.fi/handle/10250/3315</a></p> <p>Knekt, P., Lindfors, O., Ske, L. S. Ä., &amp; Virtala, E. S. A. (2013). Randomized trial on the effectiveness of long- and short-term psychotherapy on psychiatric symptoms and working ability during a 5-year follow-up. <i>Nordic journal of psychiatry</i>, 67(1), 59-68. <a href="http://www.tandfonline.com/doi/full/10.3109/08039488.2012.680910">http://www.tandfonline.com/doi/full/10.3109/08039488.2012.680910</a></p> <p>Knekt, P., Virtala, E., Härkänen, T., Vaarama, M., Lehtonen, J., &amp; Lindfors, O. (2016). The outcome of short- and long-term psychotherapy 10 years after start of treatment. <i>Psychological Medicine</i>, 1–14. <a href="http://www.journals.cambridge.org/abstract/S0033291715002718">http://www.journals.cambridge.org/abstract/S0033291715002718</a></p> <p>Lindfors, O., Knekt, P., Virtala, E., &amp; Laaksonen, M. a. (2012). The effectiveness of solution-focused therapy and short- and long-term psychodynamic psychotherapy on self-concept during a 3-year follow-up. <i>The Journal of Nervous and Mental Disease</i>, 200(11), 946–53. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23124178">http://www.ncbi.nlm.nih.gov/pubmed/23124178</a></p> <p>Lindfors, O., Knekt, P., Heinonen, E., Härkänen, T., Virtala, E., &amp; The Helsinki Psychotherapy Study Group. (2015). The effectiveness of short- and long-term psychotherapy on personality functioning during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 173, 31–8. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25462393">http://www.ncbi.nlm.nih.gov/pubmed/25462393</a></p> <p>Maljanen, T., Knekt, P., Lindfors, O., Virtala, E., Tillman, P., &amp; Härkänen, T. (2016). The cost-effectiveness of short-term and long-term psychotherapy in the treatment of depressive and anxiety disorders during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 190(July), 254–263. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22611089">http://www.ncbi.nlm.nih.gov/pubmed/22611089</a></p>	<b>Depression and/or anxiety</b>	<b>STPP</b> (Malan, 1976; Sifneos, 1978)	<p><b>1. Solution-focused therapy</b> (de Shazer et al, 1986)</p> <p><b>2. LTPP</b> (Gabbard, 2004)</p>
106.	<p><b>Knijnik, D. Z., Blanco, C., Salum, G. A., Moraes, C. U., Mombach, C., Almeida, E., ... Eizirik, C. L. (2008).</b> A pilot study of clonazepam versus psychodynamic group therapy plus clonazepam in the treatment of generalized social anxiety disorder. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 23(8), 567–74. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18774274">http://www.ncbi.nlm.nih.gov/pubmed/18774274</a></p>	<b>Social anxiety disorder (SAD)</b>	<b>Psychodynamic group therapy</b> (Malan, 1976) + <b>clonazepam</b>	<b>Clonazepam</b>
107.	<p><b>Knijnik, D. Z., Kapczinski, F., Chachamovich, E., Margis, R., &amp; Eizirik, C. L. (2004).</b> Psychodynamic group treatment for generalized social phobia. <i>Revista Brasileira de Psiquiatria (São Paulo, Brazil : 1999)</i>, 26(2), 77–81. <a href="http://www.ncbi.nlm.nih.gov/pubmed/15517">http://www.ncbi.nlm.nih.gov/pubmed/15517</a></p>	<b>Social anxiety disorder (SAD)</b>	<b>Psychodynamic group therapy</b> (Malan, 1976)	<b>Credible placebo group</b>

	<a href="#">057</a>			
108.	<b>Kompoliti, K., Wilson, B., Stebbins, G., Bernard, B., &amp; Hinson, V. (2013).</b> Immediate vs. delayed treatment of psychogenic movement disorders with short-term psychodynamic psychotherapy: Randomized clinical trial. <i>Parkinsonism &amp; related disorders</i> , 7–10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24120952">http://www.ncbi.nlm.nih.gov/pubmed/24120952</a>	<b>Psychogenic movement disorders</b>	<b>Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1980)</b>	<b>Neurological observation and support</b>
109.	<b>Kool, S., Dekker, J., &amp; Duijsens, I. (2003).</b> Efficacy of combined therapy and pharmacotherapy for depressed patients with or without personality disorders. <i>Harvard review of psychiatry</i> , 133–142. <a href="http://informahealthcare.com/doi/abs/10.1080/10673220303950">http://informahealthcare.com/doi/abs/10.1080/10673220303950</a>	<b>Depression</b>	<b>STDP (de Jonghe, 2005) + pharmacotherapy</b>	<b>Pharmacotherapy</b>
110.	<b>Kornblith, S., Rehm, L., W, O. M., &amp; Lamparski, D. M. (1983).</b> The contribution of self-reinforcement training and behavioral assignments to the efficacy of self-control therapy for depression. <i>Cogn Ther Res</i> , 7, 499-528. <a href="http://link.springer.com/10.1007/BF01172888">http://link.springer.com/10.1007/BF01172888</a>	<b>Depression</b>	<b>Psychodynamic group therapy (Wolf, 1965; Bernard &amp; Klein, 1977)</b>	<b>Self-reinforcement training and behavioral assignments (Rehm, 1977)</b>
111.	<b>Kramer, U., de Roten, Y., Perry, J. C., &amp; Despland, J.-N. (2013).</b> Change in defense mechanisms and coping patterns during the course of 2-year-long psychotherapy and psychoanalysis for recurrent depression: a pilot study of a randomized controlled trial. <i>The Journal of nervous and mental disease</i> , 201(7), 614–20. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23817160">http://www.ncbi.nlm.nih.gov/pubmed/23817160</a>	<b>Depression (recurrent)</b>	<b>Supportive-Expressive Therapy (Luborsky, 1984)</b>  All patients were also given pharmacological treatment	<b>1. Cognitive therapy (Beck, 1995)</b>  <b>2. Psychoanalysis (Robertson, 2002).</b>  <b>3. Clinical management (Csank, 2002; Novalis et al., 1993).</b>
112.	<b>Lanza, M. L., Anderson, J., Boisvert, C. M., LeBlanc, A., Fardy, M., &amp; Steel, B., et al. (2002).</b> Assaultive behavior intervention in the Veterans Administration: psychodynamic group psychotherapy compared to cognitive behavior therapy. <i>Perspect Psychiatr Care</i> , 38, 89-97. <a href="http://onlinelibrary.wiley.com/doi/10.1111/j.1744-6163.2002.tb00662.x/abstract">http://onlinelibrary.wiley.com/doi/10.1111/j.1744-6163.2002.tb00662.x/abstract</a>	<b>Male veterans with history of assault</b>	<b>Psychodynamic group psychotherapy (no specific reference)</b>	<b>Cognitive behavior group therapy (No specific reference)</b>
113.	<b>LaPonite, K., &amp; Rimm, D. C. (1980).</b> Cognitive, assertive, and insight-oriented group therapies in the treatment of reactive depression in women. <i>Psychotherapy: Theory, Research &amp; Practice</i> , 17(3), 312–321. <a href="http://psycnet.apa.org/journals/pst/17/3/312">http://psycnet.apa.org/journals/pst/17/3/312</a>	<b>Depression</b>	<b>Insight-oriented group therapy (No reference)</b>	<b>1. Cognitive group therapy (Beck, 19767)</b>  <b>2. Assertiveness training (no reference)</b>
114.	<b>Lau, M., &amp; Kristensen, E. (2007).</b> Outcome of systemic and analytic group psychotherapy for adult women with history of intrafamilial childhood sexual abuse: a randomized controlled study. <i>Acta Psychiatr Scand</i> , 116, 96-104. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17650270">http://www.ncbi.nlm.nih.gov/pubmed/17650270</a>	<b>Women with childhood sexual abuse</b>	<b>Analytic group therapy (Foulkes, 1986)</b>	<b>Systemic group therapy (Holme, 1999)</b>
115.	<b>Laurensen, E. M. P., Luyten, P., Kikkert, M. J., Westra, D., Peen, J., Soons, M. B. J., ... Dekker, J. J. M. (2018).</b> Day hospital mentalization-based treatment v. specialist treatment as usual in patients with borderline personality	<b>Borderline personality disorder</b>	<b>Day hospital mentalization-based treatment (DH-MBT; Bateman &amp; Fonagy, 2004)</b>	<b>Specialized treatment as usual (S-TAU; Van Oenen et al. 2007)</b>

	disorder: randomized controlled trial. <i>Psychological Medicine</i> , 14(1), 1–8. <a href="http://doi.org/10.1017/S0033291718000132">http://doi.org/10.1017/S0033291718000132</a>			
116.	<b>Lawson, D. M.</b> (2010). Comparing cognitive behavioral therapy and integrated cognitive behavioral therapy/psychodynamic therapy in group treatment for partner violent men. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 47(1), 122–133. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0018841">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0018841</a>	<b>Violent men</b>	<b>Integrative CBT- psychodynamic group therapy</b> (Strupp & Binder, 1984)	<b>CBT</b> (Lawson, 2001)
117.	<b>Leichsenring, F., Masuhr, O., Jaeger, U., Rabung, S., Dally, A., Dumpelmann, M., ... Streeck, U.</b> (2016). Psychoanalytic-Interactional Therapy versus Psychodynamic Therapy by Experts for Personality Disorders: A Randomized Controlled Efficacy-Effectiveness Study in Cluster B Personality Disorders. <i>Psychotherapy and Psychosomatics</i> , 71–80. <a href="http://www.karger.com/?doi=10.1159/000441731">http://www.karger.com/?doi=10.1159/000441731</a>	<b>Personality disorders (cluster B)</b>	<b>Psychoanalytic-Interactional Therapy</b> (Streeck, 2016)	<b>Psychodynamic therapy by experts</b>
118.	<b>Leichsenring, F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E.</b> (2013). Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder: A Multicenter Randomized Controlled Trial. <i>American Journal of Psychiatry</i> , 170(7), 759–767. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23680854">http://www.ncbi.nlm.nih.gov/pubmed/23680854</a>  <b>Leichsenring, F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E.</b> (2014). Long-Term Outcome of Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder. <i>American Journal of Psychiatry</i> , 171(10), 1074–1082. <a href="http://www.ncbi.nlm.nih.gov/pubmed/25016974">http://www.ncbi.nlm.nih.gov/pubmed/25016974</a>	<b>Social anxiety disorder (SAD)</b>	<b>Supportive-Expressive Therapy</b> (Luborsky, 1984)	<b>CBT</b> (Clark & Wells, 1995)
119.	<b>Leichsenring, F., Salzer, S., Jaeger, U., Kächele, H., Kreisliche, R., &amp; Leweke, F., et al.</b> (2009). Short-Term Psychodynamic Psychotherapy and Cognitive-Behavioral Therapy in Generalized Anxiety Disorder: A Randomized, Controlled Trial. <i>Am J Psychiatry</i> , 166, 875–881. <a href="http://ajp.psychiatryonline.org/article.aspx?articleID=101023&amp;amp;atab=7">http://ajp.psychiatryonline.org/article.aspx?articleID=101023&amp;amp;atab=7</a>  <b>Salzer, S., Winkelbach, C., Leweke, F., Leibing, E., &amp; Leichsenring, F.</b> (2011). Generalized Anxiety Disorder: 12-Month Follow-Up. <i>Canadian Journal of Psychiatry</i> , 56(8), 1–10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24229481">http://www.ncbi.nlm.nih.gov/pubmed/24229481</a>	<b>Generalized anxiety disorder (GAD)</b>	<b>Supportive-Expressive Therapy</b> (Luborsky, 1984)	<b>CBT</b> (Borcovec & Rucio, 2001, Brown, 1993)
120.	<b>Lemma, A., &amp; Fonagy, P.</b> (2013). Feasibility study of a psychodynamic online group intervention for depression. <i>Psychoanalytic Psychology</i> , 30(3), 367–380. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0033239">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0033239</a>	<b>Depression</b>	<b>Online Group Dynamic Interpersonal Therapy</b> (OLDIT; Fonagy & Target, 2008)	<b>Internet discussion group</b>
121.	<b>Lerner, A., Sigal, M., Bacalu, A., &amp; Gelkopf, M.</b> (1992). Short term versus long term	<b>Opiate dependence</b>	<b>Short-Term Dynamic Psychotherapy</b> (Luborsky,	<b>Long-Term Dynamic Psychotherapy</b> (Karasu,

	psychotherapy in opioid dependence: A pilot study. <i>Israeli Journal of Psychiatry and Related Sciences</i> , 29(2), 114–9. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1526756">http://www.ncbi.nlm.nih.gov/pubmed/1526756</a>		1989; Mann, 1973; Davanloo, 1978)	1989)
122.	<b>Leuzinger-Bohleber, M., Hautzinger, M., Fiedler, G., Keller, W., Bahrke, U., Kallenbach, L., ... Beutel, M. (2018).</b> Outcome of Psychoanalytic and Cognitive-Behavioural Long-Term Therapy with Chronically Depressed Patients: A Controlled Trial with Preferential and Randomized Allocation. <i>The Canadian Journal of Psychiatry</i> , 070674371878034. <a href="http://doi.org/10.1177/0706743718780340">http://doi.org/10.1177/0706743718780340</a>	<b>Depression (chronic)</b>	<b>Long-Term Psychoanalytic Psychotehrapy (LTPP;</b> Taylor, 2015)	<b>Long-term Cognitive-Behavioural Therapy (CBT;</b> Hautzinger, 2013)
123.	<b>Levy Berg, A., Sandell, R., &amp; Sandahl, C. (2009).</b> Affect-focused body psychotherapy in patients with generalized anxiety disorder: Evaluation of an integrative method. <i>Journal of Psychotherapy Integration</i> , 19(1), 67–85. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0015324">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0015324</a>	<b>Generalized anxiety disorder (GAD)</b>	<b>Affect-Focused Body Psychotherapy (Monsen &amp; Monsen, 2000)</b>	<b>TAU</b>
124.	<b>Lieberman, R. P., &amp; Eckman, T. (1981).</b> Behavior therapy vs insight-oriented therapy for repeated suicide attempters. <i>Archives of General Psychiatry</i> , 38(10), 1126–1130. <a href="http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1981.01780350060007">http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1981.01780350060007</a>	<b>Repeated suicide attempts</b>	<b>Insight-Oriented Psychotherapy</b>	<b>Behavior Therapy</b>
125.	<b>Lindauer, R. J. L., Gersons, B. P. R., Van Meijel, E. P. M., Blom, K., Carlier, I. V. E., Vrijlandt, I., &amp; Olff, M. (2005).</b> Effects of brief eclectic psychotherapy in patients with posttraumatic stress disorder: Randomized clinical trial. <i>Journal of Traumatic Stress</i> , 18(3), 205–212. <a href="http://doi.wiley.com/10.1002/its.20029">http://doi.wiley.com/10.1002/its.20029</a>	<b>PTSD</b>	<b>Brief eclectic psychotherapy (BEP;</b> Gersons et al., 2004; Manual includes both CBT and PDT components)	<b>Waitlist</b>
126.	<b>Linnet, J., &amp; Jemec, G. B. (2001).</b> Anxiety level and severity of skin condition predicts outcome of psychotherapy in atopic dermatitis patients. <i>Int J Dermatol</i> , 40, 632-636. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11737422">http://www.ncbi.nlm.nih.gov/pubmed/11737422</a>	<b>Atopic dermatitis (anxiety)</b>	<b>Brief Dynamic Psychotherapy (Malan, 1992)</b>	<b>Control (TAU?)</b>
127.	<b>Lipsitz, J. D., Gur, M., Vermes, D., Petkova, E., Cheng, J., Miller, N., ... Fyer, A. J. (2008).</b> A randomized trial of interpersonal therapy versus supportive therapy for social anxiety disorder. <i>Depression and Anxiety</i> , 25(6), 542–553. <a href="http://doi.wiley.com/10.1002/da.20364">http://doi.wiley.com/10.1002/da.20364</a>	<b>Social anxiety disorder (SAD)</b>	<b>Psychodynamic Supportive Therapy (Lipsitz et al., 2006)</b>	<b>Interpersonal Therapy (Klerman et al., 1984; Weissman et al., 2000)</b>
128.	<b>Lock, J., Grange, D. Le, &amp; Agras, W. (2010).</b> Randomized clinical trial comparing family-based treatment with adolescent-focused individual therapy for adolescents with anorexia nervosa. <i>Archives of General Psychiatry</i> , 67(10), 1025–1032. <a href="http://archpsyc.ama-assn.org/cgi/reprint/67/10/1025.pdf">http://archpsyc.ama-assn.org/cgi/reprint/67/10/1025.pdf</a>	<b>Anorexia</b>	<b>Adolescent Focused Therapy (AFT, Robin et al.)</b>	<b>Treatments Family-Based Treatment (FBT; Locke et al., 2001)</b>
129.	<b>Lorentzen, S., Ruud, T., Fjeldstad, A., &amp; Høglend, P. (2013).</b> Comparison of short- and long-term dynamic group psychotherapy: randomised clinical trial. <i>The British journal of</i>	<b>Anxiety and personality disorders</b>	<b>Long-Term Psychodynamic Group Therapy (Lorentzen, 2013)</b>	<b>Short-Term Psychodynamic Group Therapy (Lorentzen, 2013)</b>

	<p><i>psychiatry : the journal of mental science.</i>  <a href="http://www.ncbi.nlm.nih.gov/pubmed/24029539">http://www.ncbi.nlm.nih.gov/pubmed/24029539</a></p> <p>Lorentzen, S., Ruud, T., Fjeldstad, A., &amp; Høglend, P. a. (2014). Personality disorder moderates outcome in short- and long-term group analytic psychotherapy: A randomized clinical trial. <i>The British Journal of Clinical Psychology / the British Psychological Society</i>, 1–18.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/25178520">http://www.ncbi.nlm.nih.gov/pubmed/25178520</a></p> <p>Fjeldstad, A., Høglend, P., &amp; Lorentzen, S. (2015). Patterns of Change in Interpersonal Problems During and After Short-term and Long-term Psychodynamic Group Therapy: A Randomized Clinical Trial. <i>Psychotherapy Research</i>, 3307(November), 1–12.  <a href="http://www.tandfonline.com/doi/full/10.1080/10503307.2015.1102357">http://www.tandfonline.com/doi/full/10.1080/10503307.2015.1102357</a></p>			
130.	<p><b>Ludwig, G., Krenz, S., Zdrojewski, C., Bot, M., Rousselle, I., Stagno, D., ... Stiefel, F.</b> (2013). Psychodynamic interventions in cancer care I: psychometric results of a randomized controlled trial. <i>Psycho-oncology</i>, 23, 65-74.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/23983096">http://www.ncbi.nlm.nih.gov/pubmed/23983096</a></p>	<b>Mixed depression and anxiety</b> (in cancer patients)	<b>Brief Dynamic Psychotherapy</b> (Mann, 1973; Davanloo, 1994)	<b>Waitlist</b>
131.	<p><b>Maina, G., Forner, F., &amp; Bogetto, F.</b> (2005). Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. <i>Psychother Psychosom</i>, 74, 43-50.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/15627856">http://www.ncbi.nlm.nih.gov/pubmed/15627856</a></p>	<b>Depression</b> (mild to moderate)	<b>Brief Dynamic Psychotherapy</b> (BDP; Malan, 1979)	<p><b>1. Brief Supportive Psychotherapy</b> (BSP; Novalis et al., 1993)</p> <p><b>2. Waitlist</b></p>
132.	<p><b>Maina, G., Rosso, G., Crespi, C., &amp; Bogetto, F.</b> (2007). Combined brief dynamic therapy and pharmacotherapy in the treatment of major depressive disorder: a pilot study. <i>Psychotherapy and Psychosomatics</i>, 76(5), 298–305.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/17700050">http://www.ncbi.nlm.nih.gov/pubmed/17700050</a></p> <p>Maina, G., Rosso, G., &amp; Bogetto, F. (2009). Brief dynamic therapy combined with pharmacotherapy in the treatment of major depressive disorder: long-term results. <i>Journal of Affective Disorders</i>, 114(1-3), 200–7.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/18728001">http://www.ncbi.nlm.nih.gov/pubmed/18728001</a></p>	<b>Depression</b>	<b>Brief Dynamic Psychotherapy</b> (BDP; Malan, 1963; 1976) + medication	<b>Brief Supportive Psychotherapy</b> (BSP; Novalis et al., 1993) + medication
133.	<p><b>Maina, G., Rosso, G., Rigardetto, S., Chiadò Piat, S., &amp; Bogetto, F.</b> (2010). No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. <i>Psychotherapy and Psychosomatics</i>, 79(5), 295–302.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/20616624">http://www.ncbi.nlm.nih.gov/pubmed/20616624</a></p>	<b>OCD + depression</b>	<b>Brief Dynamic Psychotherapy</b> (BDP; Malan, 1979) + medication	<b>Medication only</b>
134.	<p><b>Marmar, C. R., Horowitz, M. J., Weiss, D. S., Wilner, N. R., &amp; Kaltreider, N. B.</b> (1988). A controlled trial of brief psychotherapy and mutual-help group treatment of conjugal bereavement. <i>Am J Psychiatry</i>, 145, 203-209.</p>	<b>Bereavement</b>	<b>Brief Dynamic Psychotherapy</b> (BPD; Malan, Sifneos; Mann)	<b>Mutual Self-Help Groups</b>

	<a href="http://ps.psychiatryonline.org/data/Journals/AJP/3483/203.pdf">http://ps.psychiatryonline.org/data/Journals/AJP/3483/203.pdf</a>			
135.	<b>Martini, B., Rosso, G., Chiodelli, D. F., De Cori, D., &amp; Maina, G. (2011).</b> Brief dynamic therapy combined with pharmacotherapy in the treatment of panic disorder with concurrent depressive symptoms. <i>Clinical Neuropsychiatry</i> , 8(3), 204–21. <a href="http://www.clinicalneuropsychiatry.org/pdf/04_martini.pdf">http://www.clinicalneuropsychiatry.org/pdf/04_martini.pdf</a>	<b>Panic disorder + depression</b>	<b>Brief Dynamic Psychotherapy (BDP;</b> Malan, 1963) <b>+ medication</b>	<b>Brief Supportive Psychotherapy (BSP) + medication</b>
136.	<b>McCallum, M., &amp; Piper, W. E. (1990).</b> A controlled study of effectiveness and patient suitability for short-term group psychotherapy. <i>Int J Group Psychother</i> , 40, 431-452. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2228349">http://www.ncbi.nlm.nih.gov/pubmed/2228349</a>	<b>Mixed sample</b>	<b>Short-term psychoanalytically oriented group psychotherapy (STG;</b> manual mentioned but no specific reference)	<b>Waitlist</b>
137.	<b>McLean, P., &amp; Hakstian, A. (1979).</b> Clinical depression: comparative efficacy of outpatient treatments. <i>J Consult Clin Psychol</i> , 47, 818-836. <a href="http://www.ncbi.nlm.nih.gov/pubmed/389965">http://www.ncbi.nlm.nih.gov/pubmed/389965</a>	<b>Depression</b>	<b>Short-Term Psychotherapy (Marmor,</b> 1975; Wolberg, 1967)	<b>1. Relaxation therapy (no reference)</b> <b>2. Behavior therapy (McLean, 1976)</b> <b>3. Drug therapy</b>
138.	<b>McMain, S. F., Links, P. S., Gnam, W. H., Guimond, T., Cardish, R. J., &amp; Korman, L., et al. (2009).</b> A randomized trial of dialectical behavior therapy versus general psychiatric management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1365-1374. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19755574">http://www.ncbi.nlm.nih.gov/pubmed/19755574</a>  <b>McMain, S., Guimond, T., &amp; Streiner, D. (2012).</b> Dialectical Behavior Therapy Compared With General Psychiatric Management for Borderline Personality Disorder: Clinical Outcomes and Functioning Over a 2-Year Follow-Up. <i>American Journal of Psychiatry</i> , 650–661. <a href="http://www.neuro.psychiatryonline.org/article.aspx?articleid=1148702&amp;RelatedWidgetArticles=true">http://www.neuro.psychiatryonline.org/article.aspx?articleid=1148702&amp;RelatedWidgetArticles=true</a>	<b>Borderline personality disorder</b>	<b>General psychiatric management, dynamically informed therapy (Gunderson, 1984)</b>	<b>Dialectical Behavior Therapy (DBT; Linehan et al., 1991)</b>
139.	<b>Meyer, A. E., &amp; Bolz, W. (1981).</b> The Hamburg short psychotherapy comparison experiment. Foreword and introduction. <i>Psychother Psychosom</i> , 35, 81-84. <a href="http://www.karger.com/doi/10.1159/000287484">http://www.karger.com/doi/10.1159/000287484</a>	<b>Mixed sample</b>	<b>Psychodynamic conflict-centered time-limited psychotherapy (no reference)</b>	<b>Waitlist</b>
140.	<b>Miller, L. C., Barrett, C. L., Hampe, E., &amp; Noble, H. (1972).</b> Comparison of reciprocal inhibition, psychotherapy, and waiting list control for phobic children. <i>Journal of Abnormal Psychology</i> , 79(3), 269–279. <a href="http://psycnet.apa.org/journals/abn/79/3/269L">http://psycnet.apa.org/journals/abn/79/3/269L</a>	<b>Phobia (children)</b>	<b>Play psychotherapy (Lippman, 1956)</b>	<b>Reciprocal inhibition</b>
141.	<b>Milrod, B., Chambless, D. L., Gallop, R., Busch, F. N., Schwalberg, M., McCarthy, K. S., ... Barber, J. P. (2015).</b> Psychotherapies for Panic Disorder: A Tale of Two Sites. <i>The Journal of Clinical Psychiatry</i> , 10065, 1–9. <a href="http://www.psychiatrist.com/icp/article/pages/2015/aheadofprint/14m09507.aspx">http://www.psychiatrist.com/icp/article/pages/2015/aheadofprint/14m09507.aspx</a>	<b>Panic disorder</b>	<b>Panic-Focused Psychodynamic Psychotherapy (PFPP;</b> Milrod et al, 2001)	<b>1. CBT (Craske, 2000)</b> <b>2. Applied relaxation (Öst &amp; Westling, 1995)</b>

142.	<b>Milrod, B., Leon, A. C., Busch, F., Rudden, M., Schwalberg, M., &amp; Clarkin, J., et al. (2007).</b> A randomized controlled clinical trial of psychoanalytic psychotherapy for panic disorder. <i>Am J Psychiatry, 164</i> , 265-272. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17267789">http://www.ncbi.nlm.nih.gov/pubmed/17267789</a>	<b>Panic disorder</b>	<b>Panic-Focused Psychodynamic Psychotherapy (PFPP;</b> Milrod et al, 1997)	<b>Applied relaxation</b> (Öst & Westling, 1995)
143.	<b>Molenaar, P. J., Boom, Y., Peen, J., Schoevers, R. A., Van, R., &amp; Dekker, J. J. (2011).</b> Is there a dose-effect relationship between the number of psychotherapy sessions and improvement of social functioning? <i>The British Journal of Clinical Psychology / the British Psychological Society, 50(3)</i> , 268-282. <a href="http://doi.org/10.1348/014466510X516975">http://doi.org/10.1348/014466510X516975</a>	<b>Depression</b>	<b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, Rijnierse, & Janssen, 1994)  <b>8 sessions + medication</b>	<b>Short Psychodynamic Supportive Psychotherapy</b> (de Jonghe, Rijnierse, & Janssen, 1994)  <b>16 sessions + medication</b>
144.	<b>Monsen, K., &amp; Monsen, J. T. (2000).</b> Chronic Pain and Psychodynamic Body Therapy: A Controlled Outcome Study. <i>Psychotherapy: Theory, Research, Practice, Training, 37(3)</i> , 257-269. <a href="http://psycnet.apa.org/index.cfm?fa=buy.opti onToBuy&amp;id=2000-02713-007">http://psycnet.apa.org/index.cfm?fa=buy.opti onToBuy&amp;id=2000-02713-007</a>	<b>Chronic pain</b>	<b>Psychodynamic body therapy</b> (Monsen & Monsen, 1999)	<b>TAU</b>
145.	<b>Montag, C., Haase, L., Seidel, D., Bayerl, M., Gallinat, J., Herrmann, U., &amp; Dannecker, K. (2014).</b> A Pilot RCT of Psychodynamic Group Art Therapy for Patients in Acute Psychotic Episodes: Feasibility, Impact on Symptoms and Mentalising Capacity. <i>PLoS One, 9(11)</i> , e112348. <a href="http://www.pubmedcentral.nih.gov/articleren der.fcgi?artid=4231093&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articleren der.fcgi?artid=4231093&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Schizophrenia</b>	<b>Psychodynamic art therapy</b> (Dannecker, 2010)	<b>TAU</b>
146.	<b>Mowlaie, M., Hajloo, N., Hashjin, G., Mikaeili, N., &amp; Heidari, S. (2018).</b> The Effectiveness of Empathic Dynamic Psychotherapy and Anxiety-Regulating Psychodynamic Therapy in Treating Adult Separation Anxiety Disorder. <i>Journal of Practice in Clinical Psychology, 6(3)</i> , 143-153. Retrieved from <a href="https://ejop.psychopen.eu/article/view/1528">https://ejop.psychopen.eu/article/view/1528</a>	<b>Adult Separation Anxiety Disorder (ASAD)</b>	<b>1. Empathic Dynamic Psychotherapy</b> (Seruya, 1997)  <b>2. Affect Phobia Therapy</b> (McCullough et al, 2003)	<b>Control</b> (not specified)
147.	<b>Munroe-Blum, H., &amp; Marziali, E. (1995).</b> A controlled trial of short-term group treatment for borderline personality disorder. <i>J Personal Disord, 9</i> , 190-198. <a href="http://psycnet.apa.org/psycinfo/1996-17928-001">http://psycnet.apa.org/psycinfo/1996-17928-001</a>	<b>Borderline personality disorder</b>	<b>Time-limited group therapy</b> (Marziali & Munroe-Blum, 1995)	<b>Individual psychodynamic psychotherapy</b> (Kernberg, 1975)
148.	<b>Muran, J. C., Safran, J. D., Samstag, L. W., &amp; Winston, A. (2005).</b> Evaluating an alliance-focused treatment for personality disorders. <i>Psychotherapy: Theory, Research, Practice, Training, 42(4)</i> , 532-545. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/033-3204.42.4.532">http://doi.apa.org/getdoi.cfm?doi=10.1037/033-3204.42.4.532</a>  Safran, J. D., Muran, J. C., Samstag, L. W., & Winston, A. (2005). Evaluating alliance-focused intervention for potential treatment failures: A feasibility study and descriptive analysis. <i>Psychotherapy: Theory, Research, Practice, Training, 42(4)</i> , 512-531. <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/033-3204.42.4.512">http://doi.apa.org/getdoi.cfm?doi=10.1037/033-3204.42.4.512</a>	<b>Personality disorders</b> (mixed; mainly cluster C)	<b>Brief Relational Therapy</b> (BRT; Safran & Muran, 2000)	<b>1. Short-Term Dynamic Psychotherapy</b> (STDP: Pollock, Flegenheimer, & Kaufman, 1992)  <b>2. Cognitive-behavioral therapy</b> (CBT; Turner & Muran, 1992; Beck, Freeman, & Associates, 1990)
149.	<b>Nader-Mohammadi Moghadam, M. Atef-Vahid, M. K., Asgharnejad-Farid, A. A.,</b>	<b>Social anxiety disorder (SAD)</b>	<b>Affect Phobia Therapy</b> (APT; McCullough et al.,	<b>1. Sertraline</b>



	Shabani, A., & Lavasni, F. (2015). Effectiveness of Short-term Dynamic Psychotherapy versus Sertraline in Treatment of Social Phobia. <i>Iran Journal of Psychiatry</i> , 9(2), 2–6. <a href="http://www.ncbi.nlm.nih.gov/pubmed/26288643">http://www.ncbi.nlm.nih.gov/pubmed/26288643</a>		2003)	<b>2. Waitlist</b>
150.	<b>Nijdam</b> , M. J., Gersons, B. P. R., Reitsma, J. B., de Jongh, A., & Olf, M. (2012). Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: randomised controlled trial. <i>The British Journal of Psychiatry : The Journal of Mental Science</i> , 200(3), 224–31. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22322458">http://www.ncbi.nlm.nih.gov/pubmed/22322458</a>	<b>PTSD</b>	<b>Brief eclectic psychotherapy (BEP;</b> Gersons et al, 2004; Manual includes both CBT and PDT components)	<b>Eye movement and reprocessing (EMDR;</b> De Jongh et al, 2004)
151.	<b>Nordentoft</b> , M., Øhlenschlger, J., Thorup, A., Petersen, L., Jeppesen, P., & Bertelsen, M. (2010). Deinstitutionalization revisited: A 5-year follow-up of a randomized clinical trial of hospital-based rehabilitation versus specialized assertive intervention (OPUS) versus standard treatment for patients with first-episode schizophrenia spectrum disorders. <i>Psychological Medicine</i> , 40(10), 1619–1626. <a href="http://doi.org/10.1017/S0033291709992182">http://doi.org/10.1017/S0033291709992182</a>	<b>Schizophrenia</b>	<b>Hospital-based rehabilitation</b> (included individual supportive psychodynamic psychotherapy)  Patients in all three conditions received a mix of medication, family-based psychoeducation and social skills interventions	<b>1. Assertive community training</b>  <b>2. Standard treatment</b>
152.	<b>O'Brien</b> , C. P., Hamm, K. B., Ray, B. A., Pierce, J. F., Luborsky, L., & Mintz, J. (1972). Group vs Individual Psychotherapy With Schizophrenics: A Controlled Outcome Study. <i>Archives of General Psychiatry</i> , 27(4), 474–478. <a href="http://doi.org/10.1001/archpsyc.1972.01750280042008">http://doi.org/10.1001/archpsyc.1972.01750280042008</a>	<b>Schizophrenia</b>	<b>Individual psychotherapy</b> (no reference, not clear if dynamic, but Luborsky among authors)	<b>Group psychotherapy</b> (no reference, not clear if dynamic, but Luborsky among authors)
153.	<b>Ojehagen</b> , A., Berglund, M., Appel, C. P., Andersson, K., Nilsson, B., & Skjaeris, A., et al. (1992). A randomized study of long-term outpatient treatment in alcoholics. Psychiatric treatment versus multimodal behavioural therapy, during 1 versus 2 years of treatment. <i>Alcohol Alcohol</i> , 27, 649-658. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1292438">http://www.ncbi.nlm.nih.gov/pubmed/1292438</a>	<b>Alcohol dependence</b>	<b>"Psychiatric treatment" based on psychodynamic principles</b> (Luborsky, 1984)	<b>Multimodal behavior therapy</b> (Lazarus, 1981)
154.	<b>Philips</b> , B., Wennberg, P., Konradsson, P., & Franck, J. (2018). Mentalization-Based Treatment for Concurrent Borderline Personality Disorder and Substance Use Disorder: A Randomized Controlled Feasibility Study. <i>European Addiction Research</i> , 1–8. <a href="http://doi.org/10.1159/000485564">http://doi.org/10.1159/000485564</a>	<b>Borderline personality disorder (+ substance abuse)</b>	<b>Mentalization-Based Treatment (MBT;</b> Bateman & Fonagy, 2004)  <b>+ Substance abuse treatment</b>	<b>Substance abuse treatment alone</b>
155.	<b>Pierloot</b> , R., & Vinck, J. (1978). Differential outcome of short-term dynamic psychotherapy and systematic desensitizations in the treatment of anxious outpatients: A preliminary report. <i>Psychologica Belgica</i> , 18, 87-98. <a href="http://psycnet.apa.org/psycinfo/1980-32873-001">http://psycnet.apa.org/psycinfo/1980-32873-001</a>	<b>Anxiety</b>	<b>Short-Term Dynamic Psychotherapy</b> (Malan, 1963)	<b>Systematic desensitization</b> (Wolpe, 1969)
156.	<b>Piper</b> , W. E., Azim, H. F., McCallum, M., & Joyce, A. S. (1990). Patient suitability and outcome in short-term individual psychotherapy. <i>J Consult Clin Psychol</i> , 58, 475-481. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22121">http://www.ncbi.nlm.nih.gov/pubmed/22121</a>	<b>Mixed sample</b>	<b>STDP</b> (Malan, 1976; Strupp & Binder, 1984)	<b>Waitlist</b>

	<a href="#">85</a>			
157.	<p><b>Piper, W. E., Debbane, E. G., Bienvenu, J. P., &amp; Garant, J. (1984).</b> A comparative study of four forms of psychotherapy. <i>J Consult Clin Psychol</i>, <i>52</i>, 268-279.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/6425378">http://www.ncbi.nlm.nih.gov/pubmed/6425378</a></p>	<b>Mixed sample</b>	<b>Short-term individual (STI;</b> Malan, 1976)	<p><b>1. Short-term group (STG)</b></p> <p><b>2. Long-term individual (LTI)</b></p> <p><b>3. Long-term group (LTG)</b></p> <p>Non manuals used</p>
158.	<p><b>Piper, W. E., Joyce, a S., McCallum, M., &amp; Azim, H. F. (1998).</b> Interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i>, <i>66</i>(3), 558–67.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/10224738">http://www.ncbi.nlm.nih.gov/pubmed/10224738</a></p> <p>Piper, W. E., McCallum, M., Joyce, a S., Azim, H. F., &amp; Ogrodniczuk, J. S. (1999). Follow-up findings for interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i>, <i>67</i>(2), 267–73.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/10224738">http://www.ncbi.nlm.nih.gov/pubmed/10224738</a></p>	<b>Mixed sample</b>	<b>Interpretive psychotherapy (Piper, Azim, Joyce, &amp; McCallum, 1993)</b>	<b>Supportive psychotherapy (Piper, Azim, Joyce, &amp; McCallum, 1993)</b>
159.	<p><b>Piper, W. E., McCallum, M., Joyce, A. S., Rosie, J. S., &amp; Ogrodniczuk, J. S. (2001).</b> Patient personality and time-limited group psychotherapy for complicated grief. <i>Int J Group Psychother</i>, <i>51</i>, 525-552.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/11582899">http://www.ncbi.nlm.nih.gov/pubmed/11582899</a></p> <p>Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., Weideman, R., &amp; Rosie, J. S. (2007). Group composition and group therapy for complicated grief. <i>Journal of Consulting and Clinical Psychology</i>, <i>75</i>(1), 116–25.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/17295570">http://www.ncbi.nlm.nih.gov/pubmed/17295570</a></p> <p>Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., &amp; Weideman, R. (2009). Follow-up outcome in short-term group therapy for complicated grief. <i>Group Dynamics: Theory, Research, and Practice</i>, <i>13</i>(1), 46–58.  <a href="http://doi.apa.org/getdoi.cfm?doi=10.1037/a0013830">http://doi.apa.org/getdoi.cfm?doi=10.1037/a0013830</a></p>	<b>Complicated grief</b>	<b>Interpretive group therapy (no specific reference)</b>	<b>Supportive group therapy (no specific reference)</b>
160.	<p><b>Poulsen, A. (1991).</b> Psychodynamic, time-limited group therapy in rheumatic disease--a controlled study with special reference to alexithymia. <i>Psychother Psychosom</i>, <i>56</i>, 12-23.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/1891575">http://www.ncbi.nlm.nih.gov/pubmed/1891575</a></p>	<b>Rheumatic disease</b>	<b>Time-limited group therapy (Foulkes, 1975)</b>	<b>Waitlist</b>
161.	<p><b>Poulsen, S., Lunn, S., Daniel, S. I. F., Folke, S., Mathiesen, B. B., Katznelson, H., &amp; Fairburn, C. G. (2013).</b> A Randomized Controlled Trial of Psychoanalytic Psychotherapy or Cognitive-Behavioral Therapy for Bulimia Nervosa. <i>The American journal of psychiatry</i>, 1–8.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/24275909">http://www.ncbi.nlm.nih.gov/pubmed/24275909</a></p>	<b>Bulimia</b>	<b>Psychoanalytic Psychotherapy for bulimia nervosa (Lunn &amp; Poulsen, 2012)</b>	<b>CBT (Fairburn, 2008)</b>
162.	<p><b>Qaziani, M. M., &amp; Arefi, M. (2017).</b> The effectiveness of intensive short-term dynamic psychotherapy on affection control of anxious women. <i>International Journal of Educational</i></p>	<b>Anxiety (affection control in anxious women)</b>	<b>Intensive Short-Term Dynamic Psychotherapy (no specific reference to manual)</b>	<b>No treatment waitlist</b>

	<i>and Psychological Researches</i> , 3(4). <a href="http://doi.org/10.4103/iepr.iepr">http://doi.org/10.4103/iepr.iepr</a>			
163.	<b>Ranjbar Sudejani, Y., Sharifi, K., Sayyah. S.S., &amp; Malek Mohamadi Galeh, Z. (2017)</b> Effectiveness of Intensive Short-Term Dynamic Psychotherapy (ISTDP) on Correcting Irrational Beliefs and Reducing Cognitive Avoidance of People with Obsessive Compulsive Disorder. <i>Journal of Counseling Research (article in persian)</i> , 15 (62), 140-169. <a href="http://iran-counseling.ir/journal/article-1-364-en.html">http://iran-counseling.ir/journal/article-1-364-en.html</a>	<b>OCD</b>	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Davanloo, 1990; Abbass, 2015)	<b>No treatment waitlist</b>
164.	<b>Reneses, B., Galián, M., Serrano, R., Figuera, D., Fernandez Del Moral, A., López-Ibor, J. J., ... Trujillo, M. (2013).</b> A new Time Limited Psychotherapy for BPD: Preliminary Results of a Randomized and Controlled Trial. <i>Actas españolas de psiquiatría</i> , 41(3), 139–48. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23803797">http://www.ncbi.nlm.nih.gov/pubmed/23803797</a>	<b>Borderline personality disorder</b>	<b>Representation focused Psychotherapy (PRFP;</b> (Trujillo, 2001; 2010)	<b>TAU</b>
165.	<b>Ridsdale, L., Godfrey, E., Chalder, T., Seed, P., King, M., Wallace, P., &amp; Wessely, S. (2001).</b> Chronic fatigue in general practice: is counselling as good as cognitive behaviour therapy? A UK randomised trial. <i>The British Journal of General Practice : The Journal of the Royal College of General Practitioners</i> , 51(462), 19–24. <a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1313894&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1313894&amp;tool=pmcentrez&amp;rendertype=abstract</a>	<b>Chronic fatigue</b>	<b>Counselling</b> “...all counsellors used a psychodynamic approach and all cognitive behaviour therapists used CBT” (p. 20)	<b>CBT</b>
166.	<b>O'Brien, C. P., Hamm, K. B., Ray, B. A., Pierce, J. F., Luborsky, L., &amp; Mintz, J. (1972).</b> Group vs Individual Psychotherapy With Schizophrenics: A Controlled Outcome Study. <i>Archives of General Psychiatry</i> , 27(4), 474–478. <a href="http://doi.org/10.1001/archpsyc.1972.01750280042008">http://doi.org/10.1001/archpsyc.1972.01750280042008</a>	<b>Schizophrenia</b> (20 of 24 subjects)	<b>Insight-oriented group therapy</b> (no reference)	<b>Non-insight oriented group therapy</b> (no reference)
167.	<b>Robin, A L., Siegel, P. T., &amp; Moye, a. (1995).</b> Family versus individual therapy for anorexia: impact on family conflict. <i>The International Journal of Eating Disorders</i> , 17(4), 313–22. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7620470">http://www.ncbi.nlm.nih.gov/pubmed/7620470</a>  <b>Robin, A. L., Siegel, P. T., Moye, A. W., Gilroy, M., Dennis, A. B., &amp; Sikand, A., et al. (1999).</b> A controlled comparison of family versus individual therapy for adolescents with anorexia nervosa. <i>J Am Acad Child Adolesc Psychiatry</i> , 38, 1482-1489. <a href="http://www.ncbi.nlm.nih.gov/pubmed/10596247">http://www.ncbi.nlm.nih.gov/pubmed/10596247</a>	<b>Anorexia</b>	<b>Ego-oriented individual therapy (EOIT)</b> (Robin, Siegel, Moye, Koepke, & Tice, 1994)	<b>Behavioral family systems therapy (BFST;</b> Robin et al., 1994, 1996)
168.	<b>Robinson, P., Hellier, J., Barrett, B., Barzdaitiene, D., Bateman, A., Bogaardt, A., ... Fonagy, P. (2016).</b> The NOURISHED randomised controlled trial comparing mentalisation-based treatment for eating disorders (MBT-ED) with specialist supportive clinical management (SSCM-ED) for patients with eating disorders and symptoms of borderline personality disorder. <i>Trials</i> , 17(1), 549. <a href="http://doi.org/10.1186/s13063-016-1606-8">http://doi.org/10.1186/s13063-016-1606-8</a>	<b>Borderline personality disorder + eating disorders</b>	<b>MBT-ED</b> (Bateman & Fonagy, 2009)	<b>Specialist supportive clinical management</b> (SSCM-ED; McIntosh et al., 2006)

169.	<p><b>Rosenbaum, B., Harder, S., Knudsen, P., K�ster, A., Lindhardt, A., Lajer, M., ... Winther, G. (2012).</b> Supportive Psychodynamic Psychotherapy versus Treatment as Usual for First-Episode Psychosis: Two-Year Outcome. <i>Psychiatry</i>, 75(4), 331–41. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23244011">http://www.ncbi.nlm.nih.gov/pubmed/23244011</a></p> <p>Harder, S., Koester, A., Valbak, K., &amp; Rosenbaum, B. (2014). Five-year follow-up of supportive psychodynamic psychotherapy in first-episode psychosis: long-term outcome in social functioning. <i>Psychiatry</i>, 77(2), 155–68. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24865198">http://www.ncbi.nlm.nih.gov/pubmed/24865198</a></p>	Schizophrenia	Supportive Psychodynamic Psychotherapy (SPP; Rosenbaum, 2009)	TAU
170.	<p><b>Rosser, R., Denford, J., Heslop, A., Kinston, W., Macklin, D., &amp; Minty, K., et al. (1983).</b> Breathlessness and psychiatric morbidity in chronic bronchitis and emphysema: a study of psychotherapeutic management. <i>Psychol Med</i>, 13, 93-110. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6844471">http://www.ncbi.nlm.nih.gov/pubmed/6844471</a></p>	Breathlessness and psychiatric morbidity (in patients with chronic bronchitis and emphysema)	Analytical psychotherapy (No specific reference)	<p>1. No treatment</p> <p>2. Nurse support</p>
171.	<p><b>Rosso, G., Martini, B., &amp; Maina, G. (2012).</b> Brief dynamic therapy and depression severity: A single-blind, randomized study. <i>Journal of affective disorders</i>, 1–6. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23174499">http://www.ncbi.nlm.nih.gov/pubmed/23174499</a></p>	Depression	Brief Dynamic Psychotherapy (BDP; Malan, 1976)	Brief Supportive Psychotherapy (BSP; Novalis et al., 1993)
172.	<p><b>Rossouw, T. I., &amp; Fonagy, P. (2012).</b> Mentalization-based treatment for self-harm in adolescents: a randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i>, 51(12), 1304–1313. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23200287">http://www.ncbi.nlm.nih.gov/pubmed/23200287</a></p>	Self-harm (adolescents)	Mentalization-based treatment for adolescents (MBT-A; Rossouw, unpublished)	TAU
173.	<p><b>Salminen, J. K., Karlsson, H., Hietala, J., Kajander, J., Aalto, S., &amp; Markkula, J., et al. (2008).</b> Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. <i>Psychother Psychosom</i>, 77, 351-357. <a href="http://www.ncbi.nlm.nih.gov/pubmed/18701831">http://www.ncbi.nlm.nih.gov/pubmed/18701831</a></p>	Depression	Short-term psychodynamic psychotherapy (STDP; Malan, 1976; Mann, 1973)	Fluoxetine
174.	<p><b>Salomonsson, B., &amp; Sandell, R. (2011).</b> A randomized controlled trial of mother–infant psychoanalytic treatment: I. Outcomes on self-report questionnaires and external ratings. <i>Infant Mental Health Journal</i>, 32(2), 207–231. <a href="http://onlinelibrary.wiley.com/doi/10.1002/imhj.20291/full">http://onlinelibrary.wiley.com/doi/10.1002/imhj.20291/full</a></p>	Mothers who worry about child	Mother–infant psychoanalytic treatment (MIP; Norman, 2001; 2004)	Child Health Center Care (TAU)
175.	<p><b>Salzer, S., Cropp, C., Jaeger, U., Masuhr, O., &amp; Streeck-Fischer, a. (2013).</b> Psychodynamic therapy for adolescents suffering from comorbid disorders of conduct and emotions in an in-patient setting: a randomized controlled trial. <i>Psychological medicine</i>, 1–10. <a href="http://www.ncbi.nlm.nih.gov/pubmed/24229481">http://www.ncbi.nlm.nih.gov/pubmed/24229481</a></p> <p>Salzer, S., Cropp, C., &amp; Streeck-Fischer, A. (2014). Early Intervention for Borderline Personality Disorder: Psychodynamic Therapy</p>	<p>Mixed sample of adolescents</p> <p>Adolescents with Borderline personality disorder (subsampling in secondary study)</p>	Psychoanalytic-interactive method (PiM; Streeck & Leichsenring, 2009; Leichsenring et al., 2010)	Waitlist/TAU

	in Adolescents. <i>Zeitschrift Für Psychosomatische Medizin Und Psychotherapie</i> , 60, 368–382. <a href="http://doi.org/10.13109/zptm.2014.60.4.368">http://doi.org/10.13109/zptm.2014.60.4.368</a>			
176.	<b>Salzer, S., Stefini, A., Kronmüller, K.-T., Leibing, E., Leichsenring, F., Henningsen, P., ... Steil, R. (2018).</b> Cognitive-Behavioral and Psychodynamic Therapy in Adolescents with Social Anxiety Disorder: A Multicenter Randomized Controlled Trial. <i>Psychotherapy and Psychosomatics</i> , 1–11. <a href="http://doi.org/10.1159/000488990">http://doi.org/10.1159/000488990</a>	<b>Social phobia (SAD; adolescents)</b>	<b>Supportive-Expressive Therapy</b> (Luborsky, 1984; modified for social phobia in adolescents by Horn et al, 2010)	<ol style="list-style-type: none"> <li><b>CBT</b> (Clark et al, 1995)</li> <li><b>Waitlist</b></li> </ol>
177.	<b>Sanchez, V. C., Lewinsohn, P. M., &amp; Larson, D. W. (1980).</b> Assertion training: effectiveness in the treatment of depression. <i>J Clin Psychol</i> , 36, 526-529. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7372826">http://www.ncbi.nlm.nih.gov/pubmed/7372826</a>	<b>Depression</b>	<b>"Traditional" Group Therapy</b> (No specific reference)	<b>Assertion training</b>
178.	<b>Sandahl, C., Herlitz, K., Ahlin, G., &amp; Rönnerberg, S. (1998).</b> Time-limited group psychotherapy for moderately alcohol dependent patients: A randomized controlled clinical trial. <i>Psychotherapy Research</i> , 8, 361-378. <a href="http://www.tandfonline.com/doi/abs/10.1080/10503309812331332467">http://www.tandfonline.com/doi/abs/10.1080/10503309812331332467</a>	<b>Alcohol dependence</b>	<b>Time-limited group psychotherapy</b> (Foulkes, 1964)	<b>Cognitive-behavioural group therapy</b> (Annis, 1986; Marlatt & Gordon, 1985; Sandahl and Rönnerberg, 1990)
179.	<b>Sandahl, C., Lundberg, U., Lindgren, A., Rylander, G., Herlofson, J., Nygren, A., &amp; Asberg, M. (2011).</b> Two forms of group therapy and individual treatment of work-related depression: a one-year follow-up study. <i>International journal of group psychotherapy</i> , 61(4), 539–55. <a href="http://www.ncbi.nlm.nih.gov/pubmed/21985258">http://www.ncbi.nlm.nih.gov/pubmed/21985258</a>	<b>Depression (work-related)</b>	<b>Focused psychodynamic group therapy (FGT;</b> Sandahl & Lindgren, 2003)	<ol style="list-style-type: none"> <li><b>Cognitive Group Therapy</b> (no specific reference)</li> <li><b>Comparison condition</b></li> </ol>
180.	<b>Sattel, H., Lahmann, C., Gündel, H., Guthrie, E., Kruse, J., Noll-Hussong, M., ... Henningsen, P. (2011).</b> Brief psychodynamic interpersonal psychotherapy for patients with multisomatoform disorder: randomised controlled trial. <i>The British journal of psychiatry : the journal of mental science</i> , 60–67. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22075651">http://www.ncbi.nlm.nih.gov/pubmed/22075651</a>	<b>Multisomatoform disorder</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Guthrie, 2011; Henningsen, 2011)	<b>Enhanced medical care</b>
181.	<b>Saunders, D. G. (1996).</b> Feminist-cognitive-behavioral and process-psychodynamic treatments for men who batter: interaction of abuser traits and treatment models. <i>Violence Vict</i> , 11, 393-414. <a href="http://www.ncbi.nlm.nih.gov/pubmed/9210279">http://www.ncbi.nlm.nih.gov/pubmed/9210279</a>	<b>Violent men</b>	<b>Process-psychodynamic group therapy (PPT;</b> Browne, Saunders, & Staecker, in press)	<b>Feminist-group CBT (FCBT;</b> Browne, Saunders, & Staecker, in press)
182.	<b>Schaefer, R., Kaufmann, C., Wild, B., Schellberg, D., Boelter, R., Faber, R., ... Herzog, W. (2013).</b> Specific collaborative group intervention for patients with medically unexplained symptoms in general practice: A cluster randomized controlled trial. <i>Psychotherapy and Psychosomatics</i> , 82(2), 106–119. <a href="http://doi.org/10.1159/000343652">http://doi.org/10.1159/000343652</a>	<b>Medically Unexplained Symptoms (MUS)</b>	<b>GP management + Group Interpersonal Psychodynamic Therapy</b> (Stuart & Noyes, 2006)	<b>GP management alone</b>
183.	<b>Scheidt, C. E., Waller, E., Endorf, K., Schmidt, S., König, R., Zeeck, A., ... Lacour, M. (2012).</b> Is brief psychodynamic psychotherapy in	<b>Fibromyalgia + depression</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Hobson,	<b>TAU</b>

	primary fibromyalgia syndrome with concurrent depression an effective treatment ? A randomized controlled trial. <i>General hospital psychiatry</i> . <a href="http://www.ncbi.nlm.nih.gov/pubmed/23218844">http://www.ncbi.nlm.nih.gov/pubmed/23218844</a>		1985; Goldberg et al., 1984; Guthrie, 2013)	
184.	<b>Schnyder, U., &amp; Müller, J.</b> (2011). Brief Eclectic Psychotherapy for PTSD: A Randomized Controlled Trial. <i>The Journal of Clinical ...</i> , 564–566. <a href="http://europepmc.org/abstract/MED/21527127">http://europepmc.org/abstract/MED/21527127</a>	<b>PTSD</b>	<b>Brief eclectic psychotherapy</b> (BEP; Gersons et al, 2004; Manual includes both CBT and PDT components)	<b>Minimal attention control</b>
185.	<b>Shaffer, H. J., LaSalvia, T. A., &amp; Stein, J. P.</b> (1997). Comparing Hatha yoga with dynamic group psychotherapy for enhancing methadone maintenance treatment: a randomized clinical trial. <i>Altern Ther Health Med</i> , 3, 57-66. <a href="http://www.ncbi.nlm.nih.gov/pubmed/9210777">http://www.ncbi.nlm.nih.gov/pubmed/9210777</a>	<b>Opiate addiction</b>	<b>Dynamic group psychotherapy</b>	<b>Hatha yoga therapy</b>
186.	<b>Shapiro, D. A., Barkham, M., Rees, A., Hardy, G. E., Reynolds, S., &amp; Startup, M., et al.</b> (1994). Effects of treatment duration and severity of depression on the effectiveness of cognitive-behavioral and psychodynamic-interpersonal psychotherapy. <i>J Consult Clin Psychol</i> , 62, 522-534. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8063978">http://www.ncbi.nlm.nih.gov/pubmed/8063978</a>	<b>Depression</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Hobson, 1985; Goldberg et al., 1984)	<b>Cognitive Therapy</b> (CT; Beck, Rush, Shaw, and Emery, 1979)
187.	<b>Shapiro, D., &amp; Firth, J.</b> (1987). Prescriptive v. Exploratory Psychotherapy Outcomes of the Sheffield Psychotherapy Project. <i>The British Journal of Psychiatry</i> , 151, 790-799 <a href="http://bjp.publicaciones.saludcastillayleon.es/content/151/6/790.short">http://bjp.publicaciones.saludcastillayleon.es/content/151/6/790.short</a>  Shapiro, D. a, & Firth-Cozens, J. (1990). Two-year follow-up of the Sheffield Psychotherapy Project. <i>The British journal of psychiatry</i> , 157, 389–91. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2245269">http://www.ncbi.nlm.nih.gov/pubmed/2245269</a>	<b>Depression and anxiety</b>	<b>Exploratory therapy</b> (no specific reference)	<b>Prescriptive therapy</b> (no specific reference)
188.	<b>Shaw, C. M., Margison, F. R., Guthrie, E. A., &amp; Tomenson, B.</b> (2001). Psychodynamic interpersonal therapy by inexperienced therapists in a naturalistic setting: A pilot study. <i>European Journal of Psychotherapy, Counselling, and Health</i> , 4, 87–101. <a href="http://www.tandfonline.com/doi/abs/10.1080/13642530110040091">http://www.tandfonline.com/doi/abs/10.1080/13642530110040091</a>	<b>Mixed sample</b>	<b>Psychodynamic-interpersonal psychotherapy</b> (Hobson, 1985)	<b>Waitlist</b>
189.	<b>Shefler, G., Dasberg, H., &amp; Ben-Shakhar, G.</b> (1995). A randomized controlled outcome and follow-up study of Mann's time-limited psychotherapy. <i>J Consult Clin Psychol</i> , 63, 585-593. <a href="http://psycnet.apa.org/journals/ccp/63/4/585">http://psycnet.apa.org/journals/ccp/63/4/585</a>	<b>Mixed sample</b>	<b>Time-Limited Psychotherapy</b> (TLP; Mann, 1973)	<b>Waitlist</b>
190.	<b>Simpson, S., Corney, R., Fitzgerald, P., &amp; Beecham, J.</b> (2003). A randomized controlled trial to evaluate the effectiveness and costeffectiveness of psychodynamic counselling for general practice patients with chronic depression. <i>Psychol Med</i> , 33, 229-239. <a href="http://www.ncbi.nlm.nih.gov/pubmed/12622">http://www.ncbi.nlm.nih.gov/pubmed/12622</a>	<b>Depression (chronic)</b>	<b>Psychodynamic counselling</b> (Burton, 1998)	<b>General practice</b>

	<a href="#">302</a>			
191.	<b>Sjodin, I., Svedlund, J., Ottosson, J. O., &amp; Dotevall, G. (1986).</b> Controlled study of psychotherapy in chronic peptic ulcer disease. <i>Psychosomatics</i> , 27, 187-91, 195. <a href="http://www.psychosomaticsjournal.com/article/S0033-3182(86)72711-4/abstract">http://www.psychosomaticsjournal.com/article/S0033-3182(86)72711-4/abstract</a>	<b>Peptic ulcer disease</b>	<b>Short-Term Dynamic Therapy (Malan, 1976)</b>	<b>Medication</b>
192.	<b>Sleed, M., Baradon, T., &amp; Fonagy, P. (2013).</b> New Beginnings for mothers and babies in prison: A cluster randomized controlled trial. <i>Attachment &amp; Human Development</i> , 15(4), 349–67. <a href="http://www.ncbi.nlm.nih.gov/pubmed/23550526">http://www.ncbi.nlm.nih.gov/pubmed/23550526</a>	<b>High risk mothers (and their babies) in prison</b>	<b>Attachment-based intervention (Baradon, 2006)</b>	<b>Standard health and social care</b>
193.	<b>Sloane, R. B., Staples, F. R., Cristol, A. H., Yorkston, N. J., &amp; Whipple, K. (1975).</b> Short-term analytically oriented psychotherapy versus behavior therapy. <i>American Journal of Psychiatry</i> , 132, 373-377. <a href="http://www.ncbi.nlm.nih.gov/pubmed/1119588">http://www.ncbi.nlm.nih.gov/pubmed/1119588</a>	<b>Anxiety or personality disorder</b>	<b>Short-Term Analytically Oriented Psychotherapy (no specific reference)</b>	<b>1. Waitlist</b> <b>2. Behavior Therapy (no specific reference)</b>
194.	<b>Smyrniotis, K. X., &amp; Kirkby, R. J. (1993).</b> Long-term comparison of brief versus unlimited psychodynamic treatments with children and their parents. <i>J Consult Clin Psychol</i> , 61, 1020-1027. <a href="http://www.ncbi.nlm.nih.gov/pubmed/8113479">http://www.ncbi.nlm.nih.gov/pubmed/8113479</a>	<b>Children with mixed problems</b>	<b>Time-limited brief psychotherapy (Smyrniotis, Kirkby, Smyrniotis, &amp; Picone, 1987)</b>	<b>Time-unlimited psychotherapy (Klein, 1975; Segal, 1978)</b>
195.	<b>Snyder, D. K., Wills, R. M., &amp; Grady-Fletcher, a. (1991).</b> Long-term effectiveness of behavioral versus insight-oriented marital therapy: a 4-year follow-up study. <i>Journal of Consulting and Clinical Psychology</i> , 59(1), 138–41. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2002129">http://www.ncbi.nlm.nih.gov/pubmed/2002129</a>	<b>Couples with relational problems</b>	<b>Insight-Oriented Marital Therapy (IOMT; no specific reference)</b>	<b>Behavioral Marital Therapy (BMT; no specific reference)</b>
196.	<b>Soares, M. C., Mondin, T. C., Silva, G. D. G. da, Barbosa, L. P., Molina, M. L., Jansen, K., ... Silva, R. A. da. (2018).</b> Comparison of Clinical Significance of Cognitive-Behavioral Therapy and Psychodynamic Therapy for Major Depressive Disorder. <i>The Journal of Nervous and Mental Disease</i> , 00(00), 1. <a href="http://doi.org/10.1097/NMD.0000000000000872">http://doi.org/10.1097/NMD.0000000000000872</a>	<b>Major Depression (MDD)</b>	<b>Supportive-Expressive Therapy (SET; Luborsky, 1984)</b>	<b>Cognitive-Behavioral Therapy (CBT; Beck et al, 1997)</b>
197.	<b>Sørensen, P., Birket-Smith, M., Wattar, U., Buemann, I., &amp; Salkovskis, P. (2010).</b> A randomized clinical trial of cognitive behavioural therapy versus short-term psychodynamic psychotherapy versus no intervention for patients with hypochondriasis. <i>Psychol Med</i> , 1-11. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20380779">http://www.ncbi.nlm.nih.gov/pubmed/20380779</a>	<b>Hypochondriasis</b>	<b>Short-Term Psychodynamic Psychotherapy (No specific reference)</b>	<b>1. Cognitive-behavioural Therapy (CBT; Salkovskis, 1989; Warwick, 1989; Salkovskis et al. 2003)</b> <b>2. Waitlist</b>
198.	<b>Spiess, K., Sachs, G., Pietschmann, P., &amp; Prager, R. (1995).</b> A program to reduce onset distress in unselected type I diabetic patients: Effects on psychological variables and metabolic control. <i>European Journal of Endocrinology</i> . Retrieved from <a href="http://ovidsp.ovid.com/ovidweb.cgi?T=JS&amp;PAGE=reference&amp;D=emed3&amp;NEWS=N&amp;AN=1995149187">http://ovidsp.ovid.com/ovidweb.cgi?T=JS&amp;PAGE=reference&amp;D=emed3&amp;NEWS=N&amp;AN=1995149187</a>	<b>Type I diabetic patients</b>	<b>Group PDT program (no specific manual or reference) + treatment as usual</b>	<b>Treatment as usual</b>

199.	<p><b>Stanton, A. H., Gunderson, J. G., Knapp, P. H., Frank, A. F., Vancelli, M. L., &amp; Schnitzer, R., et al. (1984).</b> Effects of psychotherapy in schizophrenia: I. Design and implementation of a controlled study. <i>Schizophr Bull</i>, 10, 520-563.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/6151245">http://www.ncbi.nlm.nih.gov/pubmed/6151245</a></p>	Schizophrenia (non-chronic)	Expressive, insight-oriented psychotherapy (EIO; Stanton, in press)	Reality-adaptive, supportive therapy (RAS; May, 1968)
200.	<p><b>Stefini A, Salzer S, Reich G, Horn H, Winkelmann K, Bents H, Rutz U, Frost U, Boetticher Av, Ruhl U, Specht N, Kronmüller K-T. (2017).</b> Cognitive-Behavioral and Psychodynamic Therapy in Female Adolescents With Bulimia Nervosa: A Randomized Controlled Trial. <i>Journal of the American Academy of Child &amp; Adolescent Psychiatry</i>, 56(4), 329-335.  <a href="http://doi.org/10.1016/j.jaac.2017.01.019">http://doi.org/10.1016/j.jaac.2017.01.019</a></p>	Bulimia (in female adolescents)	Focal psychodynamic therapy for bulimia (Reich et al, 1998; 2014)	CBT (Bents, 2007; 2010; Fairburn, 2008)
201.	<p><b>Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattheß, H., ... Kruse, J. (2016).</b> Resource activation for treating post-traumatic stress disorder, comorbid symptoms and impaired functioning: a randomized controlled trial in Cambodia. <i>Psychological Medicine</i>, 1–12.  <a href="http://doi.org/10.1017/S0033291716002592">http://doi.org/10.1017/S0033291716002592</a></p> <p><b>Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattes, H., ... Kruse, J. (2016).</b> Treating post-traumatic stress disorder by resource activation in Cambodia. <i>World Psychiatry</i>, 15(2), 182–183.  <a href="http://doi.org/10.1002/wps.20312">http://doi.org/10.1002/wps.20312</a></p>	PTSD	Resource-Oriented Trauma Therapy (ROTATE; Wöller & Mattheß, 2016)	Waitlist
202.	<p><b>Suchman, N. E., Decoste, C. L., Castiglioni, N., McMahon, T. J., Rounsaville, B. J., &amp; Mayes, L. (2011).</b> The Mothers and Toddlers Program, an attachment-based parenting intervention for substance using women: Post- treatment results from a randomized clinical pilot. <i>Attachment &amp; Human Development</i>, 12(5), 483–504.  <a href="http://doi.org/10.1080/14616734.2010.501983">http://doi.org/10.1080/14616734.2010.501983</a></p>	Mothers with alcohol and drug addiction	The Mothers and Toddlers Program (MTP)	Parent Education (PE)
203.	<p><b>Suchman, N. E., Decoste, C. L., McMahon, T. J., Dalton, R., Mayes, L. C., &amp; Borelli, J. (2017).</b> Mothering from the Inside Out: Results of a second randomized clinical trial testing a mentalization-based intervention for mothers in addiction treatment. <i>Development and Psychopathology</i>, 29(2), 617–636.  <a href="http://doi.org/10.1017/S0954579417000220">http://doi.org/10.1017/S0954579417000220</a></p>	Mothers with histories of alcohol and drug addiction	Mothering from the Inside Out (MIO; Suchman & Bers, 2015)	Parent Education (PE)
204.	<p><b>Svartberg, M., Seltzer, M. H., &amp; Stiles, T. C. (1998).</b> The effects of common and specific factors in short-term anxiety-provoking psychotherapy: a pilot process-outcome study. <i>J Nerv Ment Dis</i>, 186, 691-696.  <a href="http://journals.lww.com/jonmd/Abstract/1998/11000/The_Effects_of_Common_and_Specific_Factors_in.5.aspx">http://journals.lww.com/jonmd/Abstract/1998/11000/The_Effects_of_Common_and_Specific_Factors_in.5.aspx</a></p>	Mixed sample	Short-term anxiety-provoking psychotherapy (STAPP; Sifneos, 1992)	Non-directive Psychotherapy (NDP; utveckledes specifikt för studien)
205.	<p><b>Svartberg, M., Stiles, T. C., &amp; Seltzer, M. H. (2004).</b> Randomized, controlled trial of the effectiveness of short-term dynamic psychotherapy and cognitive therapy for cluster C personality disorders. <i>Am J Psychiatry</i>, 161, 810-817.  <a href="http://www.ncbi.nlm.nih.gov/pubmed/15121">http://www.ncbi.nlm.nih.gov/pubmed/15121</a></p>	Personality disorder (Cluster C)	Short-Term Dynamic Psychotherapy (STDP; McCullough Vaillant, 1997)	Cognitive Therapy (CT; Beck & Freeman, 1990)



	<a href="#">645</a>			
206.	<p>Svedlund, J., Sjodin, I., Ottosson, J. O., &amp; Dotevall, G. (1983). Controlled study of psychotherapy in irritable bowel syndrome. <i>Lancet</i>, 2, 589-592.</p> <p><a href="http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3240791&amp;tool=pmcentrez&amp;rendertype=abstract">http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3240791&amp;tool=pmcentrez&amp;rendertype=abstract</a></p>	IBS	Brief Dynamic Psychotherapy (Malan, 1979)	Treatment as usual (TAU) (medical)
207.	<p>Szapocznik, J., Rio, A., Murray, E., Cohen, R., Scopetta, M., &amp; Rivas-Vazquez, A., et al. (1989). Structural family versus psychodynamic child therapy for problematic Hispanic boys. <i>J. Consult. Clin. Psychol.</i>, 57, 571-578.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/2794177">http://www.ncbi.nlm.nih.gov/pubmed/2794177</a></p>	Behavioral problems among boys	Individual psychodynamic child therapy (IPCT) (Adams, 1974; Cooper & Wanerman, 1977)	<p>1. Structural Family Therapy (SFT; Minuchin &amp; Fishman, 1981; Minuchin, Montalvo, Guerney, Rosman, &amp; Schuman, 1967)</p> <p>2. Recreational control condition</p>
208.	<p>Tasca, G., Ritchie, K., Conrad, G., Balfour, L., Gayton, J., &amp; Lybanon, V., et al. (2006). Attachment scales predict outcome in a randomized controlled trial of two group therapies for binge eating disorder: An aptitude by treatment interaction. <i>Psychotherapy Research</i>, 16, 106-121.</p> <p><a href="http://www.tandfonline.com/doi/abs/10.1080/10503300500090928">http://www.tandfonline.com/doi/abs/10.1080/10503300500090928</a></p>	Binge eating disorder	Group psychodynamic interpersonal psychotherapy (Tasca et al., 2002; Strupp & Binder, 1984; Malan 1979)	Group cognitive-behavior therapy (GCBT; Wilfley et al., 1996)
209.	<p>Tasca, G. A., Koszycki, D., Brugnera, A., Chyurlia, L., Hammond, N., Francis, K., ... Balfour, L. (2018). Testing a stepped care model for binge-eating disorder: a two-step randomized controlled trial. <i>Psychological Medicine</i>.</p> <p><a href="http://doi.org/10.1017/S0033291718001277">http://doi.org/10.1017/S0033291718001277</a></p>	Binge eating disorder	Group psychodynamic interpersonal psychotherapy (Tasca et al., 2002; Strupp & Binder, 1984; Malan 1979)	No treatment control
210.	<p>Thompson, L. W., Gallagher, D., &amp; Breckenridge, J. S. (1987). Comparative effectiveness of psychotherapies for depressed elders. <i>J Consult Clin Psychol</i>, 55, 385-390.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/3597953">http://www.ncbi.nlm.nih.gov/pubmed/3597953</a></p>	Depression (among elderly)	Brief psychodynamic therapy (Horowitz & Kaltreider, 1979)	<p>1. Cognitive Therapy (CT; Beck, Rush, Shaw, &amp; Emery, 1979)</p> <p>2. Behavioral Therapy (BT; Gallagher &amp; Thompson, 1981; Lewinsohn, 1974)</p> <p>3. Delayed-treatment control</p>
211.	<p>Thyme, K. E., Sundin, E. C., Stahlberg, G., Lindstrom, B., Eklof, H., &amp; Wiberg, B., et al. (2007). The outcome of short-term psychodynamic art therapy compared to short-term psychodynamic verbal therapy for depressed women. <i>Psychoanalytic Psychotherapy</i>, 21, 250-264.</p> <p><a href="http://www.tandfonline.com/doi/abs/10.1080/02668730701535610">http://www.tandfonline.com/doi/abs/10.1080/02668730701535610</a></p>	Depression (among women)	Short-Term Psychodynamic Verbal Therapy (Mann, 1973)	Art psychotherapy (no reference)
212.	<p>Trowell, J., Joffe, I., Campbell, J., Clemente, C., Almqvist, F., &amp; Soininen, M., et al. (2007). Childhood depression: a place for psychotherapy. An outcome study comparing individual psychodynamic psychotherapy and family therapy. <i>Eur Child Adolesc Psychiatry</i>, 16, 157-167.</p> <p><a href="http://www.ncbi.nlm.nih.gov/pubmed/17200793">http://www.ncbi.nlm.nih.gov/pubmed/17200793</a></p> <p>Garoff, F. F., Heinonen, K., Pesonen, A.-K., &amp; Almqvist, F. (2012). Depressed youth:</p>	Depression (children)	Focused Individual Psychotherapy (Malan & Osmimo, 1992; Davanloo, 1978)	Systems Integrative Family Therapy (Byung-Hall, 1995; Will, 1985)

	<p>treatment outcome and changes in family functioning in individual and family therapy. <i>Journal of Family Therapy</i>, 34(1), 4–23. <a href="http://doi.wiley.com/10.1111/j.1467-6427.2011.00541.x">http://doi.wiley.com/10.1111/j.1467-6427.2011.00541.x</a></p>			
213.	<p><b>Trowell, J., Kolvin, I., Weeramanthri, T., Sadowski, H., Berelowitz, M., &amp; Glaser, D., et al. (2002).</b> Psychotherapy for sexually abused girls: psychopathological outcome findings and patterns of change. <i>Br J Psychiatry</i>, 180, 234-247. <a href="http://www.ncbi.nlm.nih.gov/pubmed/11872516">http://www.ncbi.nlm.nih.gov/pubmed/11872516</a></p>	<b>Trauma</b> (sexually abused girls)	<b>Individual therapy</b> (Unclear with no reference)	<b>Psychopedagogical /analytic group therapy</b>
214.	<p><b>Town, J. M., Abbass, A., Stride, C. &amp; Bernier, D. (2017)</b> A randomised controlled trial of Intensive Short-Term Dynamic Psychotherapy for treatment resistant depression: the Halifax Depression Study, <i>Journal of Affective Disorders</i>, in Press <a href="http://dx.doi.org/10.1016/j.jad.2017.02.035">http://dx.doi.org/10.1016/j.jad.2017.02.035</a></p>	<b>Depression</b> (treatment resistant)	<b>Intensive Short-Term Dynamic Psychotherapy</b> (ISTDP; Abbass, 2015)	<b>TAU</b> (Canadian depression guidelines)
215.	<p><b>Vinnars, B., Barber, J. P., Norén, K., Gallop, R., &amp; Weinryb, R. M. (2005).</b> Manualized supportive-expressive psychotherapy versus nonmanualized community-delivered psychodynamic therapy for patients with personality disorders: bridging efficacy and effectiveness. <i>Am J Psychiatry</i>, 162, 1933-1940. <a href="http://www.ncbi.nlm.nih.gov/pubmed/16199841">http://www.ncbi.nlm.nih.gov/pubmed/16199841</a></p> <p>Vinnars, B., Gallop, R., &amp; Barber, J. P. (2009). Do Personality Problems Improve During Psychodynamic Supportive-Expressive Psychotherapy? Secondary Outcome Results From a Randomized Controlled Trial for Psychiatric Outpatients with Personality Disorders. <i>Psychotherapy</i>, 46, 362–375. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20161588">http://www.ncbi.nlm.nih.gov/pubmed/20161588</a></p>	<b>Personality disorder</b> (mixed)	<b>Supportive-Expressive Psychotherapy</b> (SE; Luborsky, 1984)	<b>Non-manualized PDT</b>
216.	<p><b>Vitriol, V. G., Ballesteros, S. T., Florenzano, R. U., Weil, K. P., &amp; Benadof, D. F. (2009).</b> Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. <i>Psychiatr Serv</i>, 60, 936-942. <a href="http://www.ncbi.nlm.nih.gov/pubmed/19564224">http://www.ncbi.nlm.nih.gov/pubmed/19564224</a></p>	<b>Depression</b> (+ childhood trauma)	<b>Brief psychodynamic psychotherapy</b> (Kudler et al, 2003; Davies et al, 1994)	<b>TAU</b>
217.	<p><b>Watzke, B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., &amp; Schulz, H. (2010).</b> Effectiveness of systematic treatment selection for psychodynamic and cognitive-behavioural therapy: randomised controlled trial in routine mental healthcare. <i>The British Journal of Psychiatry : The Journal of Mental Science</i>, 197(2), 96–105. <a href="http://www.ncbi.nlm.nih.gov/pubmed/20679260">http://www.ncbi.nlm.nih.gov/pubmed/20679260</a></p> <p>Watzke, B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., &amp; Schulz, H. (2012). Longer term outcome of cognitive-behavioural and psychodynamic psychotherapy in routine mental health care: Randomised controlled trial. <i>Behaviour research and therapy</i>, 50(9), 580–587.</p>	<b>Mixed sample</b>	<b>Psychodynamic brief group therapy</b> (Heigl-Evers et al., 1997)	<b>Cognitive-behavioural group therapy</b> (Fiedler, 2001)

	<a href="http://www.ncbi.nlm.nih.gov/pubmed/22750189">http://www.ncbi.nlm.nih.gov/pubmed/22750189</a>			
218.	<b>Wiborg, I. M., &amp; Dahl, A. A. (1996).</b> Does brief dynamic psychotherapy reduce the relapse rate of panic disorder? <i>Arch Gen Psychiatry</i> , 53, 689-694. <a href="http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1996.01830080041008">http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1996.01830080041008</a>	<b>Panic disorder</b>	<b>Brief Dynamic Psychotherapy (BDP;</b> Davanloo, 1978; Malan 1976; 1979; Strupp & Binder, 1984) <b>+ Clomipramin</b>	<b>Clomipramin alone</b>
219.	<b>Wiltink, J., Dippel, A., Szczepanski, M., Thiede, R., Alt, C., &amp; Beutel, M. E., et al. (2007).</b> Long-term weight loss maintenance after inpatient psychotherapy of severely obese patients based on a randomized study: predictors and maintaining factors of health behavior. <i>J Psychosom Res</i> , 62, 691-698. <a href="http://www.ncbi.nlm.nih.gov/pubmed/17540227">http://www.ncbi.nlm.nih.gov/pubmed/17540227</a>	<b>Severe obesity</b>	<b>Psychodynamic inpatient treatment</b> (no manual but reference to: Levine & Luborsky, 1981)	<b>Behavioral treatment</b> (no specific reference)
220.	<b>Wiltink, J., Ruckes, C., Hoyer, J., Leichsenring, F., Joraschky, P., Leweke, F., ... Beutel, M. E. (2017).</b> Transfer of manualized Short Term Psychodynamic Psychotherapy (STPP) for social anxiety disorder into clinical practice: results from a cluster-randomised controlled trial. <i>BMC Psychiatry</i> , 17(1), 92. <a href="http://doi.org/10.1186/s12888-017-1257-7">http://doi.org/10.1186/s12888-017-1257-7</a>	<b>Social Phobia</b>	<b>Manualized Short-Term Psychodynamic Psychotherapy</b> (based on Supportive-Expressive Therapy; Luborsky, 1984)	<b>Non-manualized Short-Term Psychodynamic Psychotherapy</b> (tauSTPP; no manual)
221.	<b>Winston, A., Pollack, J., McCullough, L., Flegenheimer, W., Kestenbaum, R., &amp; Trujillo, M., et al. (1991).</b> Brief psychotherapy of personality disorders. <i>J Nerv Ment Dis</i> , 179, 188-193. <a href="http://psycnet.apa.org/psycinfo/1991-25198-001">http://psycnet.apa.org/psycinfo/1991-25198-001</a>  Winston, A, Laikin, M., Pollack, J., Samstag, L. W., McCullough, L., & Muran, J. C. (1994). Short-term psychotherapy of personality disorders. <i>The American journal of psychiatry</i> , 151(2), 190-4. <a href="http://www.ncbi.nlm.nih.gov/pubmed/22217220">http://www.ncbi.nlm.nih.gov/pubmed/22217220</a>	<b>Personality disorders</b> (mixed)	<b>Short-Term Dynamic Therapy</b> (STDP; Davanloo, 1980; Also reference to Mann, Malan, and Sifneos)	<b>Brief Adaptive Psychotherapy</b> (BAP; Pollack et al., 1991)
222.	<b>Woody, G. E., Luborsky, L., McLellan, A. T., O'Brien, C. P., Beck, A. T., &amp; Blaine, J., et al. (1983).</b> Psychotherapy for opiate addicts. Does it help? <i>Arch Gen Psychiatry</i> , 40, 639-645. <a href="http://www.ncbi.nlm.nih.gov/pubmed/6847332">http://www.ncbi.nlm.nih.gov/pubmed/6847332</a>  Woody, G. E., Luborsky, L., Mcllellan, A. T., & O'Brien, C. P. (1990). Corrections and Revised analyses for Psychotherapy in Methadone Patients. <i>Arch Gen Psychiatry</i> , 47, 788-789. <a href="http://www.ncbi.nlm.nih.gov/pubmed/2378551">http://www.ncbi.nlm.nih.gov/pubmed/2378551</a>	<b>Opiate dependence</b>	<b>Supportive-Expressive Psychotherapy</b> (SE; Luborsky, 1977; unpublished manual)	<b>Cognitive-Behavior Therapy</b> (CBT; Beck, unpublished)  <b>Drug Counseling</b> (DC; Stockdale & Hargrove, unpublished material, 1977)
223.	<b>Woody, G. E., McLellan, A. T., Luborsky, L., &amp; O'Brien, C. P. (1995).</b> Psychotherapy in community methadone programs: A validation study. <i>American Journal of Psychiatry</i> , 152, 1302-1308. <a href="http://www.ncbi.nlm.nih.gov/pubmed/7653685">http://www.ncbi.nlm.nih.gov/pubmed/7653685</a>	<b>Opiate dependence</b>	<b>Supportive-Expressive Psychotherapy</b> (SE; Luborsky, 1984)	<b>Drug counseling</b>
224.	<b>Zipfel, S., Wild, B., Groß, G., Friederich, H., Teufel, M., Schellberg, D., ... Zwaan, M. De. (2013).</b> Focal psychodynamic therapy,	<b>Anorexia</b>	<b>Focal psychodynamic therapy</b> (FPT; Schauenburg et al., 2009)	<b>1. Enhanced cognitive behavior therapy</b> (CBT-E; Fairburn, 2008)

	cognitive behaviour therapy, and optimised treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised controlled trial. <i>The Lancet</i> , 383, 127–137. <a href="http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)61746-8/abstract">http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)61746-8/abstract</a>			<b>2. Optimized TAU</b>
225.	<b>Zwerenz, R., Becker, J., Gerzymisch, K., Siepmann, M., Holme, M., Kiwus, U., ... Beutel, M. E. (2017).</b> Evaluation of a transdiagnostic psychodynamic online intervention to support return to work: A randomized controlled trial. <i>Plos One</i> , 12(5), e0176513. <a href="http://doi.org/10.1371/journal.pone.0176513">http://doi.org/10.1371/journal.pone.0176513</a>	<b>Work-related stress</b>	<b>Online intervention based on Supportive-Expressive Psychotherapy (SE;</b> Luborsky, 1984)	<b>Online control group</b>
226.	<b>Zwerenz, R., Becker, J., Johansson, R., Frederick, R. J., Andersson, G., &amp; Beutel, M. E. (2017).</b> Transdiagnostic, Psychodynamic Web-Based Self-Help Intervention Following Inpatient Psychotherapy: Results of a Feasibility Study and Randomized Controlled Trial. <i>JMIR Mental Health</i> , 4(4), e41. <a href="http://doi.org/10.2196/mental.7889">http://doi.org/10.2196/mental.7889</a>	<b>Mixed sample</b>	<b>IPDT (Affect-focused;</b> based on Frederick, 2009)	<b>Waitlist</b>
227.	<b>Zwerenz, R., Becker, J., Knickenberg, R. J., Siepmann, M., Hagen, K., &amp; Beutel, M. E. (2017).</b> Online Self-Help as an Add-On to Inpatient Psychotherapy: Efficacy of a New Blended Treatment Approach. <i>Psychotherapy and Psychosomatics</i> , 341–350. <a href="http://doi.org/10.1159/000481177">http://doi.org/10.1159/000481177</a>	<b>Depression (inpatients)</b>	<b>Inpatient Psychodynamic Psychotherapy (program of different interventions, no manual referenced) + Internet intervention (Deprexis)</b>	<b>Inpatient Psychodynamic Psychotherapy (program of different interventions, no manual referenced) + Internet control intervention (psychoeducation)</b>